

Product Bulletin



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Chaparral™

EPA Reg. No. 62719-597

2(ee) Recommendation†

Control of Susceptible Weeds Pre or Post Planting of Certain Annual and Perennial Grasses

ATTENTION

† This recommendation is permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA.

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Read and follow all applicable directions for use, precautions, and limitations on the product label attached to the container for Chaparral™ herbicide.

Directions for Use

Chaparral controls susceptible broadleaf plants both pre and postemergence. See the Chaparral main label for a list of susceptible weeds.

PRECAUTIONS

Postemergence applications on newly emerged tall fescue, or newly emerged or established annual ryegrass, timothy, sorghum-sudangrass, teff, crabgrass, and pearl millet are not recommended. Unacceptable injury or death of these grasses may occur with Chaparral applications.

Pre-plant applications for control of susceptible emerged weeds

Apply Chaparral at a maximum of 2.0 oz/acre with a non-ionic surfactant to control susceptible weeds no sooner than 30 days prior to planting tall fescue, orchardgrass, timothy, and annual ryegrass. Chaparral can also be used at 2.0 oz/acre prior to planting sorghum-sudangrass, teff, crabgrass, and pearl millet, but a minimum 45-day-interval must be observed between the application and planting.

Post planting applications for control of susceptible emerged weeds

Apply Chaparral at a maximum of 2.0 oz/acre with a non-ionic surfactant to orchardgrass after seedling plants have at least begun tillering (multiple stems from single crown).

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

R371-023

Issued: 06/02/2017

Initial printing