

Product Bulletin



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

GrazonNext[®] HL

EPA Reg. No. 62719-628

2(ee) Recommendation[†]

Control of Susceptible Weeds Pre or Post Planting of Certain Annual and Perennial Grasses

ATTENTION

[†] This recommendation is permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA.

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Read and follow all applicable directions for use, precautions, and limitations on the product label attached to the container for GrazonNext[®] HL herbicide.

Directions for Use

GrazonNext HL controls susceptible broadleaf plants both pre and postemergence. See the GrazonNext HL main label for a list of susceptible weeds.

PRECAUTIONS

Postemergence applications on newly emerged or established sorghum-sudangrass, teff, crabgrass, and pearl millet are not recommended. Unacceptable injury or death of these grasses may occur with GrazonNext HL applications.

Pre-plant applications for control of susceptible emerged weeds

Apply GrazonNext HL at a maximum of 1.5 pts/acre with a non-ionic surfactant to control susceptible weeds no sooner than 15 days prior to planting tall fescue, orchardgrass, timothy, and annual ryegrass. GrazonNext HL can also be used at 1.5 pts/acre prior to planting sorghum-sudangrass, teff, crabgrass, and pearl millet, but a minimum 30-day-interval must be observed between the application and planting.

Post planting applications for control of susceptible emerged weeds

Apply GrazonNext HL at a maximum of 1.5 pts/acre with a non-ionic surfactant to tall fescue, orchardgrass, timothy, and annual ryegrass after seedling plants have at least reached the V3 stage (three collared leaves).

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

R392-025

Issued: 05/26/2017

Initial printing