

---

## SUPPLEMENTAL LABELING

---

**Syngenta Crop Protection, LLC**  
P. O. Box 18300  
Greensboro, North Carolina 27419-8300  
**SCP 1324A-S7 1016**

GROUP	3	11	FUNGICIDES
-------	---	----	------------

### **Quilt Xcel® Fungicide**

Broad spectrum fungicide for control of plant diseases

**This supplemental label expires on October 21, 2019 and must not be used or distributed after this date.**

#### Active Ingredients:

Azoxystrobin.....	13.5%
Propiconazole .....	11.7%
Other Ingredients:	74.8%
Total:	100.0%

Contains 1.02 lb ai propiconazole and 1.18 lb ai azoxystrobin per gallon

Quilt Xcel Fungicide is formulated as a suspo-emulsion (SE).

**KEEP OUT OF REACH OF CHILDREN.**

## **WARNING/AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

### **EPA Reg. 100-1324**

**All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using Quilt Xcel Fungicide as permitted according to this Supplemental Labeling, read and follow all applicable directions, restrictions, and precautions on the EPA-registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and/or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.**

**syngenta**

## DIRECTIONS FOR USE

<b>Rotational Crops</b>	<b>Planting Time From Last Quilt Xcel Application</b>
Brassica, leafy greens (subgroup 5B) Bulb crops Carrots Celery (and other leaf petiole crops - subgroup 4B) Cereals (wheat, barley, triticale) Corn (field, seed, popcorn, and sweet) Dill Grasses grown for seed (Cool season grasses only) Mint Oats Peanuts Quinoa Radish Rice Rye Sorghum Soybeans Strawberries Sugar beets Wild rice Watercress	0 days
Buckwheat Millet	12 Months
Alfalfa (if propiconazole rate does not exceed 0.22 lb ai/acre/season)	75 days
All Other Crops Intended for Food and Feed	105 days

Crop	Target Diseases	Use Rate fl oz product/A	Remarks
<b>Grasses grown For Seed</b> (Cool season grasses only)	Ergot Stem Diseases Powdery Mildew ( <i>Erysiphe graminis</i> ) Rusts ( <i>Puccinia</i> spp.)	14 - 26	<p>Apply Quilt Xcel when powdery mildew infections, and/or rust pustules are noticeable and increasing in number in late spring or early summer. To maximize control of severe rust pressure, apply 26 fl oz/A (except bluegrass apply 14 fl oz/A) and make applications at 14-day intervals until the seed is mature. For bluegrass, it is important to begin application early in the growing season.</p> <p>Make no more than 2 sequential applications of a Group 11 fungicide before alternating to another product with a different mode of action than Group 11 fungicides.</p>

**Application:** Quilt Xcel is most effective when applied and allowed to dry before a rainfall. For best results, sufficient coverage is very important. Apply Quilt Xcel in a minimum of 20 gal of water per acre for ground or in a minimum of 10 gal of water per acre for aerial. Quilt Xcel may be applied by ground, air, or chemigation.

**Specific Use Restrictions:**

- 1) Do not feed hay cut within 20 days of the last application.
- 2) Do not graze treated areas within 140 days of the last application.
- 3) Do not apply more than 86.0 fl oz/A/season of Quilt Xcel.
- 4) Do not apply more than 0.90 lb ai propiconazole-containing products/A/season.
- 5) Do not apply more than 0.8 lb ai azoxystrobin-containing products/A/season.
- 6) Do not apply within 20 days of harvest (20-day PHI) of seed.
- 7) Do not apply to bermudagrass grown for seed.

Crop	Target Diseases	Use Rate fl oz product/A	Remarks
Mint  Peppermint Spearmint	Powdery Mildew ( <i>Erysiphe</i> spp.) Rust ( <i>Puccinia menthae</i> )	10.5 - 14	Begin applications when the plants are 2-4 inches high or when conditions become favorable for disease development. Make a second application 14 days after the first application.

**Application:** Quilt Xcel may be applied by aerial, ground (minimum of 20 gal/A) or chemigation.

**Specific Use Restrictions:**

- 1) Do not apply more than 28 fl oz/A of Quilt Xcel per crop.
- 2) Do not apply more than 0.22 lb ai of propiconazole-containing products/A/season.
- 3) Do not apply more than 0.75 lb ai of azoxystrobin-containing products/A/season.
- 4) Do not apply within 7 days of harvest (7-day PHI).

Quilt Xcel® and the Syngenta logo are trademarks of a Syngenta Group Company

©2017 Syngenta

Quilt Xcel SCP 1324A-S7 1016 – ep – 5-4-17