



Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

SUPPLEMENTAL LABEL

*This supplemental label **expires on 03/02/2019**
and must not be used or distributed after this
date.*

LUNA® EXPERIENCE

FOR USE ON: Almond; Brassica, leafy greens, subgroup 5B; Cherry subgroup 12-12A; Fruit, small vine climbing, except fuzzy kiwifruit, subgroup 13-07F; Hops, dried cones; Legume Vegetables (except cowpea; dried peas and soybeans); Melon subgroup 9A; Nut, tree, group 14-12; Okra; Onion, bulb, subgroup 3-07A; Onion, green, subgroup 3-07B; Peach subgroup 12-12B; Pistachio; Plum subgroup 12-12C; Squash/Cucumber subgroup 9B; Sunflower subgroup 20B.

ACTIVE INGREDIENT:

FLUOPYRAM*..... 17.6%

TEBUCONAZOLE*..... 17.6%

OTHER INGREDIENTS:..... 64.8%

TOTAL: 100.0%

Contains 1.67 lbs FLUOPYRAM and 1.67 lbs TEBUCONAZOLE per gallon

*(CAS Numbers 658066-35-4 and 107534-96-3)

EPA Reg. No. 264-1091

EPA Est. No.

SUSPENSION CONCENTRATE

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This Supplemental Label and the entire LUNA EXPERIENCE container label, EPA Reg. No 264-1091, must be in possession of the user at the time of application.
- Read the label affixed to the container for LUNA EXPERIENCE before applying.
- Use of LUNA EXPERIENCE according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for LUNA EXPERIENCE. Refer to the LUNA EXPERIENCE container label for additional instructions and restrictions.

Aerial Application Prohibited in New York State.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during **the restricted-entry interval (REI) of 12 hours for all crops except when performing cane tying, turning, or girdling on grapes. The REI is 5 days for treated grapes when conducting cane tying, turning, or girdling on grapes. The REI is 12 hours for all other activities in grapes.**

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls over long-sleeved shirt and long pants, socks and shoes, and chemical-resistant gloves made of any waterproof material such as natural rubber \geq 14 mils.

Ground Application

For optimum disease control, apply in sufficient water to ensure thorough coverage of foliage, bloom, and fruit.

Aerial Application

For aerial application equipment, a minimum of **10** gallons of water per acre for tree crops and **2** gallons of water per acre for field and vegetable crops is required. Aerial Application Prohibited in New York State.

ROTATIONAL CROP RESTRICTIONS

Areas treated with this product can be replanted immediately following harvest with any crop for which both a FLUOPYRAM and a TEBUCONAZOLE tolerance exist. This includes: Almond; Barley; Brassica, leafy greens, subgroup 5B; Cherry subgroup 12-12A; Corn, field, grain; Corn, pop, grain; Corn, sweet kernel plus cob with husks removed; Cotton; Fruit, pome, group 11-10; Fruit, small vine climbing, except fuzzy kiwifruit, subgroup 13-07F; Hops, dried cones; Melon subgroup 9A; Nut, tree, group 14-12; Okra; Onion, bulb, subgroup 3-07A; Onion, green, subgroup 3-07B; Peach subgroup 12-12B; Peanut; Pepper/Eggplant subgroup 8-10B; Pecan; Pistachio; Plum subgroup 12-12C; Soybean; Sunflower subgroup 20B; Tomato subgroup 8-10A; Squash/Cucumber subgroup 9B; Wheat.

The following crops can be planted into treated areas 120 days after the last application LUNA EXPERIENCE: Alfalfa; Artichoke, globe; Berry, low growing, except cranberry, subgroup 13-07G; Bushberry subgroup 13-07B; Caneberry subgroup 13-07A Carrot; Dill seed; Fruit, citrus, group 10-10; Ginseng; Grain, cereal, group 15 (except corn and rice) which include: buckwheat, millet (pearl and proso), oats, rye, sorghum, teosinte, and triticale; Herb subgroup 19A; Leafy greens subgroup 4A; Leaf petioles subgroup 4B (except watercress);

Rapeseed subgroup 20A; Sugarcane (in Region 3); Sugarbeet, root; Vegetable, legume, Pea and bean, succulent shelled, subgroup 6B; Vegetable, root, except sugarbeet, subgroup 1B; Vegetable, tuberous and corm, subgroup 1C.

Do not rotate to crops other than those listed above.

USE DIRECTIONS FOR SPECIFIC CROPS

Almond		
Disease Control	Application Rate	Product Instructions
Brown rot blossom blight <i>(Monilinia laxa)</i> <i>(Monilinia fructicola)</i> Shot hole <i>(Wilsonomyces carpophilus)</i> Powdery mildew <i>(Sphaerotheca pannosa)</i> <i>(Podosphaera tridactyla)</i> Jacket rot* Green fruit rot <i>(Botrytis cinerea)</i> Alternaria leaf spot* <i>(Alternaria alternata)</i> Anthracnose* <i>(Colletotrichum acutatum)</i> Scab* <i>(Cladosporium carpophilum)</i> Rust* <i>(Tranzschelia discolor)</i>	6.0 to 17 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Disease Suppression	Application Rate	Product Instructions
Hull rot* <i>(Rhizopus spp.)</i> <i>(Monilinia spp.)</i>	6.0 to 17 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Restrictions: <ul style="list-style-type: none"> Do not apply more than 34 fl oz of LUNA EXPERIENCE per acre per year. Apply using ground, aerial, or chemigation equipment. Regardless of formulation or method of application, do not apply more than 0.446 lbs Fluopyram or 0.900 lbs tebuconazole per acre per year, including soil and foliar uses. Do not apply LUNA EXPERIENCE within 35 days of harvest. 		

- Do not cut cover crops in treated areas for feed, or allow livestock to graze treated areas.
- To limit the potential for development of disease resistance to these fungicide classes, do not make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.

*Not for use in CA without a supplemental label.

Brassica, Leafy greens, subgroup 5B

Leafy Greens subgroup: Broccoli Raab; Cabbage, Chinese (Bok Choy); Collards; Kale; Mizuna; Mustard Greens; Mustard Spinach; Rape Greens.

Disease Control	Application Rate	Product Instructions
Powdery mildew* <i>(Erysiphe cruciferarum)</i> <i>(Erysiphe polygoni)</i> Cercospora leaf spot* <i>(Cercospora brassicicola)</i> Alternaria blight / Black spot* <i>(Alternaria spp.)</i>	6.0 to 8.6 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Botrytis gray mold <i>(Botrytis cinerea)</i> Sclerotinia stem rot <i>(Sclerotinia sclerotiorum)</i> <i>(Sclerotinia minor)</i>	8.6 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 14-day interval. When disease pressure is severe, use the shorter intervals.
Disease Suppression	Application Rate	Product Instructions
Rhizoctonia bottom rot <i>(Rhizoctonia solani)</i>	8.6 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 14-day interval. When disease pressure is severe, use the shorter intervals.

Restrictions:

- Do not apply more than 34 fl oz of LUNA EXPERIENCE per acre per year.
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, do not apply more than 0.446 lbs Fluopyram or 0.450 lbs tebuconazole per acre per year, including soil and foliar uses.
- Do not apply LUNA EXPERIENCE within 7 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, do not make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.
- Application to Turnip greens is limited to east of the Rockies.

*Not for use in CA without a supplemental label.

Cherry subgroup 12-12A; Peach subgroup 12-12B; and Plum subgroup 12-12C

Cherry subgroup: Capulin; Cherry, black; Cherry, Nanking; Cherry, sweet; Cherry, tart; cultivars, varieties, and/or hybrids of these.

Peach subgroup: Peach; Nectarine; cultivars, varieties, and/or hybrids of these.

Plum subgroup: Apricot; Apricot, Japanese; Jujube, Chinese; Plum; Plum, American; Plum, beach; Plum, Canada; Plum, cherry; Plum, Chickasaw; Plum, Damson; Plum, Japanese; Plum, Klamath; Plumcot; Plum, prune; Sloe; cultivars, varieties, and/or hybrids of these.

Disease Control	Application Rate	Product Instructions
Brown rot blossom blight Fruit rot (<i>Monilinia</i> spp.) Powdery mildew Rusty spot (<i>Podosphaera</i> spp.) (<i>Sphaerotheca pannosa</i>) Shot hole* (<i>Wilsonomyces carpophilus</i>) Jacket rot, Green fruit rot* (<i>Botrytis cinerea</i>) Peach leaf curl* (<i>Taphrina</i> spp.)	6.0 to 10 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Scab (<i>Cladosporium carpophilum</i>) Rust (<i>Tranzschelia discolor</i>)	8.0 to 10 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.

Restrictions:

- Do not apply more than 34 fl oz of LUNA EXPERIENCE per acre per year.
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, do not apply more than 0.446 lbs Fluopyram or 1.35 lbs tebuconazole per acre per year, including soil and foliar uses.
- May be applied the day of harvest. To limit the potential for development of disease resistance to these fungicide classes, do not make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.

*Not for use in CA without a supplemental label.

Fruit, small vine climbing, except fuzzy kiwifruit, subgroup 13-07F

Amur River Grape; Gooseberry; Grape; Kiwifruit, Hardy; Maypop; Schisandra Berry; cultivars varieties, and/or hybrids of these.

Disease Control	Application Rate	Product Instructions
Powdery mildew (<i>Uncinula necator</i>)	6.0 to 8.6 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 12- to 21-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Botrytis bunch rot / Gray mold (<i>Botrytis cinerea</i>) Black rot (<i>Guignardia bidwellii</i>) Sour rot* (<i>various pathogens</i>)	8.0 to 8.6 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 12- to 21-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Disease Suppression	Application Rate	Product Instructions
Phomopsis cane and leaf spot (<i>Phomopsis viticola</i>)	8.6 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 12- to 21-day interval. When disease pressure is severe, use the shorter intervals.

Restrictions:

- Do not apply more than 34 fl oz of LUNA EXPERIENCE per acre per year.
- Apply using ground, aerial, or chemigation equipment. Regardless of formulation or method of application, do not apply more than 0.446 lbs Fluopyram or 0.900 lbs tebuconazole per acre per year, including soil and foliar uses.
- Do not apply LUNA EXPERIENCE within 14 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, do not make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.

*Not for use in CA without a supplemental label.

Hops, dried cones		
Disease Control	Application Rate	Product Instructions
Powdery mildew (<i>Sphaerotheca humuli</i>) (<i>Sphaerotheca macularis</i>)	8.0 to 17 fl oz/acre	<p>Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 14-day interval.</p> <p>Make applications in sufficient water volume (just to the point of runoff). Thorough coverage is needed for disease control. Apply increasing product rate as plant size increases.</p> <p>8.0 fl oz with 15 - 30 gals./acre 10.0 fl oz with 31 - 60 gals./acre 12.0 fl oz with 61 - 90 gals./acre 14.0 fl oz with 91 - 200 gals./acre</p>
Disease Suppression	Application Rate	Product Instructions
Gray mold (<i>Botrytis</i> spp.)	17 fl oz/acre	<p>Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 14-day interval.</p>
Restrictions: <ul style="list-style-type: none"> Do not apply more than 34 fl oz of LUNA EXPERIENCE per acre per year. Apply using ground, aerial, or chemigation equipment. Regardless of formulation or method of application, do not apply more than 0.446 lbs Fluopyram or 0.900 lbs tebuconazole per acre per year, including soil and foliar uses. Do not apply LUNA EXPERIENCE within 14 days of harvest. To limit the potential for development of disease resistance to these fungicide classes, do not make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group. 		

Legume Vegetables (except cowpea; dried peas and soybeans) *

Edible-Podded Legume Vegetables subgroup A: Bean (*Phaseolus* spp.) (Includes Runner Bean, Snap Bean, Wax Bean); Bean (*Vigna* spp.) (Includes Asparagus Bean, Chinese Longbean, Moth Bean, Yardlong Bean); Jackbean; Pea (*Pisum* spp.) (Includes Dwarf Pea, Edible-Pod Pea, Snow Pea, Sugar Snap Pea); Pigeon Pea; Sword Bean.

Dried Shelled Bean (Except Soybean and Dried Shelled Pea) subgroup C: Dried Cultivars Of Bean (*Lupinus* spp.) (includes Grain Lupin, Sweet Lupin, White Lupin, and White Sweet Lupin); (*Phaseolus* spp.) (includes Field Bean, Kidney Bean, Lima Bean (Dry), Navy Bean, Pinto Bean; Tepary Bean; Bean (*Vigna* spp.) (includes Adzuki Bean, Blackeyed Pea, Catjang, Cowpea, Crowder Pea, Moth Bean, Mung Bean, Rice Bean, Southern Pea, Urd Bean); Broad Bean (Dry); Chickpea; Guar; Lablab Bean; Lentil; Pea (*Pisum* spp.) (includes Field Pea); Pigeon Pea.

Disease Control	Application Rate	Product Instructions
Powdery mildew (<i>Erysiphe graminis</i>) Septoria leaf and glume blotch (<i>Septoria</i> spp.) Rusts (<i>Puccinia</i> spp.)	8.0 to 12.8 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 10- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.

Restrictions:

- Do not apply more than 25.6 fl oz of LUNA EXPERIENCE per acre per year.
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, do not apply more than 0.446 lbs Fluopyram or 0.672 lbs tebuconazole per acre per year, including soil and foliar uses.
- Do not apply LUNA EXPERIENCE within 14 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, do not make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.
- Do not allow livestock to graze treated area for 14 days and do not harvest for food or feed within 14 days of application.

*Not for use in CA without a supplemental label.

Melon subgroup 9A and Squash/Cucumber subgroup 9B:**Melon subgroup:** Citron Melon; Muskmelon; Watermelon.**Squash/Cucumber subgroup:** Chayote (Fruit); Chinese Waxgourd; Cucumber; Gherkin; Gourd, Edible; *Momordica* spp.; Pumpkin; Squash, Summer; Squash, Winter.

Disease Control	Application Rate	Product Instructions
Powdery mildew (<i>Sphaerotheca fuliginea</i> / <i>Podosphaera xanthii</i>) (<i>Erysiphe cichoracearum</i>) Alternaria leaf spot (<i>Alternaria cucumerina</i>)	6.0 to 17 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 10- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Gummy stem blight* (<i>Didymella bryonae</i>) Belly rot* (<i>Rhizoctonia solani</i>) Anthracnose* (<i>Colletotrichum</i> spp.)	10.0 to 17 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 10- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.

Note: A mild yellowing on leaf margins is sometimes seen following application of LUNA EXPERIENCE in cucurbits.**Restrictions:**

- Do not apply more than 34 fl oz of LUNA EXPERIENCE per acre per year.
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, do not apply more than 0.446 lbs Fluopyram or 0.691 lbs tebuconazole per acre per year, including soil and foliar uses.
- Do not apply LUNA EXPERIENCE within 7 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, do not make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.

*Not for use in CA without a supplemental label.

Nut, tree, group 14-12**(See separate use directions for: Almonds, and Pistachio)**

African Nut-Tree; Beechnut; Brazil Nut; Brazilian Pine; Bunya; Bur Oak; Butternut; Cajou Nut; Candlenut; Cashew; Chestnut; Chinquapin; Coconut; Coquito Nut; Dika Nut; Ginkgo; Guiana Chestnut; Hazelnut; Heartnut; Hickory Nut; Japanese Horse-Chestnut; Macadamia Nut; Mongongo Nut; Monkey-Pot; Monkey Puzzle Nut; Okari Nut; Pachira Nut; Peach Palm Nut; Pequi; Pili Nut; Pine Nut; Sapucaia Nut; Tropical Almond; Walnut, Black; Walnut, English; Yellowhorn; Cultivars, varieties, and/or hybrids of these.

Disease Control	Application Rate	Product Instructions
Botryosphaeria panicle and shoot blight <i>(Botryosphaeria dothidea)</i> Brown rot blossom blight <i>(Monilinia laxa)</i> <i>(Monilinia fructicola)</i> Alternaria leaf and fruit spot <i>(Alternaria alternata)</i> Alternaria late blight <i>(Alternaria alternata)</i> Scab <i>(Cladosporium carpophilum)</i> <i>(Cladosporium caryigenum)</i> Shothole <i>(Wilsonomyces carpophilus)</i> Anthracnose <i>(Colletotrichum acutatum)</i> <i>(Glomerella cingulata)</i>	10.0 to 17 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 14- to 21-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Disease Suppression	Application Rate	Product Instructions
Eastern filbert blight <i>(Anisogramma anomala)</i>	10.0 to 17 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 7- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.

Restrictions:

- Do not apply more than 34 fl oz of LUNA EXPERIENCE per acre per year.
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, do not apply more than 0.446 lbs Fluopyram or 0.900 lbs tebuconazole per acre per year, including soil and foliar uses.
- Do not apply LUNA EXPERIENCE within 35 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, do not make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.
- Do not cut cover crops in treated areas for feed or allow livestock to graze treated areas.

Okra

Disease Control	Application Rate	Product Instructions
Cercospora leaf spot (<i>Cercospora</i> spp.)	12.8 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 14-day interval.

Restrictions:

- Do not apply more than 34 fl oz of LUNA EXPERIENCE per acre per year.
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, do not apply more than 0.446 lbs Fluopyram or 0.675 lbs tebuconazole per acre per year including soil and foliar uses.
- Do not apply LUNA EXPERIENCE within 3 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, do not make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.

Onion, bulb, subgroup 3-07A and Onion, green, subgroup 3-07B

Onion, bulb subgroup: Daylily, Bulb; Fritillaria, Bulb; Garlic, Bulb; Garlic, Great-Headed, Bulb; Garlic, Serpent, Bulb; Lily, Bulb; Onion, Bulb; Onion, Chinese, Bulb; Onion, Pearl; Onion, Potato, Bulb; Shallot, Bulb; cultivars, varieties, and/or hybrids of these.

Onion, green subgroup: Chive, Fresh Leaves; Chive, Chinese, Fresh Leaves; Elegans Hosta; Fritillaria, Leaves; Kurrat; Lady's Leek; Leek; Leek, Wild; Onion, Beltsville Bunching; Onion, Fresh; Onion, Green; Onion, Macrostem; Onion, Tree, Tops; Onion, Welsh, Tops; Shallot, Fresh Leaves; cultivars, varieties, and/or hybrids of these.

Disease Control	Application Rate	Product Instructions
Botrytis leaf blight, neck rot <i>(Botrytis squamosa)</i> <i>(Botrytis allii)</i> Purple blotch <i>(Alternaria porri)</i> Rust <i>(Puccinia allii)</i> <i>(Puccinia porri)</i>	8.0 to 12.8 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 10- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Stemphylium leaf blight / stalk rot <i>(Stemphylium vesicarium)</i> Cladosporium leaf blotch <i>(Cladosporium allii)</i>	12.8 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 10- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Disease Suppression	Application Rate	Product Instructions
White rot <i>(Sclerotium cepivorum)</i>	12.8 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 10- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.

Restrictions:

- Do not apply more than 25.6 fl oz of LUNA EXPERIENCE per acre per year.
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, do not apply more than 0.446 lbs Fluopyram or 0.335 lbs tebuconazole per acre per year, including soil and foliar uses.
- Do not apply LUNA EXPERIENCE within 7 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, do not make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.

Pistachio		
Disease Control	Application Rate	Product Instructions
Botryosphaeria panicle and shoot blight <i>(Botryosphaeria dothidea)</i> Septoria leaf spot <i>(Septoria pistaciarum)</i> Blossom and shoot blight <i>(Botrytis cinerea)</i> Alternaria late blight <i>(Alternaria alternata)</i>	6.4 to 17 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 10- to 14-day interval. When disease pressure is severe, use the higher rates and/or shorter intervals.
Restrictions: <ul style="list-style-type: none"> Do not apply more than 34 fl oz of LUNA EXPERIENCE per acre per year. Apply using ground, aerial, or chemigation equipment. Regardless of formulation or method of application, do not apply more than 0.446 lbs Fluopyram or 0.900 lbs tebuconazole per acre per year, including soil and foliar uses. Do not apply LUNA EXPERIENCE within 35 days of harvest. To limit the potential for development of disease resistance to these fungicide classes, do not make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group. Do not cut cover crops in treated areas for feed, or allow livestock to graze treated areas. 		

Sunflower subgroup 20B

Calendula; Castor Oil Plant; Chinese Tallowtree; Euphorbia; Evening Primrose; Jojoba; Niger Seed; Rose Hip; Safflower; Stokes Aster; Sunflower; Tallowwood; Tea Oil Plant; Vernonia; cultivars, varieties, and/or hybrids of these.

Disease Control	Application Rate	Product Instructions
Rust (<i>Puccinia helianthi</i>) Alternaria leaf spot (<i>Alternaria</i> spp.) Sclerotinia wilt and head rot (<i>Sclerotinia sclerotiorum</i>) Powdery mildew (<i>Erysiphe cichoracearum</i>)	9.0 to 12.8 fl oz/acre	Apply at the critical timings for disease control. Refer to University and/or extension guidelines for best application timings. Continue as needed on a 14-day interval. When disease pressure is severe, use the higher rates.

Restrictions:

- Do not apply more than 34 fl oz of LUNA EXPERIENCE per acre per year.
- Apply using ground, aerial, or chemigation equipment.
- Regardless of formulation or method of application, do not apply more than 0.446 lbs Fluopyram or 0.450 lbs tebuconazole per acre per year, including soil and foliar uses.
- Do not apply LUNA EXPERIENCE within 50 days of harvest.
- To limit the potential for development of disease resistance to these fungicide classes, do not make more than 2 sequential applications of LUNA EXPERIENCE or any Group 7 or Group 3 containing fungicide before rotating with a fungicide from a different Group.

For **MEDICAL** and **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577
For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

SAL on 03/02/2016AV3

LUNA is a Registered trademark of Bayer.