

Supplemental Labeling



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Grasp[®] SC

EPA Reg. No. 62719-500

This supplemental label expires **March 13, 2021** and must not be used or distributed after this date.

For Pre-emergence weed control in rice in the states of Arkansas, Florida, Louisiana, Mississippi, Missouri and Texas.

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Grasp[®] SC before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Grasp SC according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Grasp SC.

Application Information

Use Rate

Apply 2 oz Grasp[®] SC per acre (0.031 lb ai/ac) in rice as a pre-emergence application. One post-emergence application of Grasp[®] SC may be made provided the total amount of Grasp[®] SC applied during one year does not exceed 5.6 ounces per acre in the first and ratoon crop

Application Timing

Grasp[®] SC herbicide may be applied to rice as a pre-emergence or post-emergence application (in drill seeded rice) or rice pegging at 1 leaf stage with no exposed roots (in water seeded rice) up to 60 days before harvest. Within this application window, application timing is dependent on cultural practices and optimum timing for weed species present. (See Application Rates and Weeds Controlled table.) Do not apply if crop or weeds are under drought stress.

Tank Mixtures

Grasp[®] SC may be used alone or tankmixed with one or more registered products according to the specific tank mixing instructions in this label and respective product labels. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Pre-emergence Application Rates and Weeds Controlled¹

Arkansas, Florida, Louisiana, Mississippi, Missouri, and Texas

| Weeds Controlled Common name (<i>scientific name</i>) | Application Rate 2.0 fl oz/acre |
|---|--|
| eclipta (<i>Eclipta alba</i>) Indian/northern jointvetch (<i>Aeschynomene</i> spp.) rice flatsedge (<i>Cyperus iria</i>) smartweed spp, annual (<i>Polygonum spp</i>) | |
| Barnyardgrass (<i>Echinochloa-crus-galli</i>) junglerice (<i>Echinochloa-colona</i>) pigweed (<i>Amaranthus spp</i>) | |

¹ Grasp may not control known ALS resistant weed biotypes.

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

R194-015
EPA accepted 02/14/18
Initial Printing