

**RESTRICTED USE PESTICIDE
TOXIC TO FISH AND AQUATIC ORGANISMS**
FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER
THEIR DIRECT SUPERVISION AND ONLY FOR THE USES COVERED BY THE CERTIFIED
APPLICATOR'S CERTIFICATION.



Fanfare[®] 2EC Insecticide/Miticide
EPA REG. NO. 66222-99

FIFRA §2(ee) RECOMMENDATION

**FIFRA SECTION 2(ee) RECOMMENDATION FOR USE IN THE STATES OF
ALABAMA, FLORIDA, GEORGIA, NORTH CAROLINA, SOUTH CAROLINA,
TENNESSEE, AND VIRGINIA**

**For Control of Kudzu Bug (*Megacopta cribraria*) in Soybeans and Legume
Vegetables**

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the US EPA.
- Observe all applicable precautions, restrictions, limitations, and instructions on the EPA-approved label.
- This labeling must be in the possession of the user at the time of pesticide application.

APPLICATION INFORMATION

Fanfare[®] 2EC can be applied for the control of kudzu bug a rate of 6.4 fluid ounces per acre.

LB AI/A	FL OZ/A
0.100	6.4

For the full list of Legume Vegetables, see the label for FANFARE[®] 2EC.

Clemson Cooperative Extension recommends Fanfare 2EC for controlling kudzu bugs on soybeans and comments:

Under certain conditions, kudzu bug can cause economic losses if not controlled. Insecticides should be timed for optimum control of the immature stage. Trigger initial application when immature are first detected in sweep-net samples at approximately 1 nymph/sweep. An application at R3/R4 will likely prevent this generation from completing development on soybean. Thereafter, use 2 bugs/sweep if populations build again.

Refer to the label for Fanfare[®] 2EC for additional use directions, restrictions, rotational crop intervals and precautions.

5/25/12
EPA 062411



3120 Highwoods Blvd, Suite 100, Raleigh NC 27604
T: 919.256.9305 ♦ F: 919.256.9308