

STAYLO™

DEPOSITION AID / DRIFT RETARDANT

ACTIVE INGREDIENTS:

Polyvinyl Polymer (Polyacrylamide)	30%
Inert Ingredients	70%
TOTAL	100%

All ingredients are exempt from tolerance requirements under 40 CFR 180.1001 (d).

KEEP OUT OF REACH OF CHILDREN WARNING

Hazards to humans and domestic animals.

Causes skin irritation. Do not get in eyes, on skin or clothing. Causes moderate eye irritation.

GENERAL INFORMATION

STAYLO is an effective, easy-to-use adjuvant for deposition improvement and drift retardation in spraying operations. **STAYLO** is compatible with most water soluble and wettable powder pesticides, desiccant and cotton defoliants when applied by aerial application or standard ground equipment.

STAYLO when used as directed in accordance with sound agricultural practices, is in compliance with Federal Food and Drug Administration Regulations governing the cropland use of adjuvants and is exempt from tolerance requirements under Title 40, Code of Federal Regulations 180.1001 (d).

DIRECTIONS FOR USE

Important: Keep container closed in storage and do not allow water to come in contact with contents until added to the spray solution.

Step 1. Select correct dosage from chart below.

Step 2. Fill mix tank with water, pesticide and other additives and begin to agitate.

Step 3. SHAKE **STAYLO** BOTTLE WELL BEFORE USING.

Step 4. For best mixing results, either (1) inject **STAYLO** liquid into the suction side of the feeder or recirculating pump to obtain adequate dispersion of the polymer throughout the pesticide solution, or (2) slowly add **STAYLO** liquid to the rapidly agitating tank mix in the area of highest turbulence by pouring a very thin stream.

Step 5. Continue to agitate mix tank for at least 2 minutes before spraying.

Step 6. Follow all cleanup precautions on label of pesticide used. Follow governmental procedures for disposal of pesticide spray solution.

NOTE: The degree of drift hazard varies with the type of pesticide, application conditions, and vegetation near the sprayed area. Consult your local agricultural advisor.

Remember, pesticide drift is no accident. Common sense and sound application technology must be followed when spraying pesticides. **STAYLO** will retard, but not totally eliminate drift.

DOSAGE CHART

Spray Pressure	Nozzle Orientation	STAYLO *Dosage
aerial applied below 45 psi	straight back 45° angle back	2-6 oz. 4-8 oz.
ground application low (below 30 psi)	flat fan, flooding off-center	1-4 oz. 3-5 oz.
medium (below 30-50 psi)	flat fan, flooding off-center spray guns	3-4 oz. 4-6 oz. 5-8 oz.
high (over 50 psi)	spray guns	8-32 oz.

* Ounces per 100 gallons of spray solution.

For home and garden sprayers, use approximately 1 Tablespoon of **STAYLO** per 5 gallons of spray solution.

FIRST AID

If on Skin: Wash with soap and water. If irritation develops, consult a physician. Remove and launder contaminated clothing before reuse.

If in Eyes: Flush with water for at least 15 minutes. Seek medical attention if irritation persists.

If Inhaled: Remove to fresh air, treat symptomatically.

STORAGE AND DISPOSAL

PROTECT FROM FREEZING. Store above 32°F. Do not reuse empty container. Dispose of container in a safe manner in accordance with federal, state and local regulations.

WARRANTY

Seller warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to the seller, and buyer assumes the risk or any such use.



Manufactured For:
TRIANGLE CHEMICAL COMPANY
MACON, GEORGIA

Manufactured For:
CARDINAL CHEMICALS, INC.
KINSTON, NC