



# Supplemental Label

## ASSAIL® 30SG INSECTICIDE

EPA Reg. No. 8033-36-70506

**This supplemental label expires on December 31, 2019, and must not be used or distributed after this date.**

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Follow all applicable directions, restrictions and precautions on the federally-registered label.

This label must be in the possession of the user at the time of pesticide application. Read the label affixed to the container for ASSAIL 30SG Insecticide before applying. Use of ASSAIL 30SG Insecticide according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container of ASSAIL 30SG Insecticide.

### GRAPES and other Climbing Vine Small Fruits (except Fuzzy Kiwifruit) (within Crop Sub-Group 13-07F)

#### SPRAY VOLUME FOR GRAPES and OTHER CLIMBING VINE SMALL FRUITS (EXCEPT KIWI FRUIT) (within Crop Sub-Group 13-07F):

Apply in a minimum finished spray volume of 5 gallons per acre by air or 20 gallons per acre by ground. Ground applications are recommended for optimal control.

SITE	PEST	DOSAGE PER ACRE		USE DIRECTIONS
		Pounds active	Ounces Assail 30SG Insecticide	
<b>GRAPES and other Climbing Vine Small Fruits (except Fuzzy Kiwifruit) (within Crop Sub-Group 13-07F):</b> Amur river grape; gooseberry, hardy kiwifruit, Maypop, schisandra berry, and cultivars, varieties, and/or hybrids of these.	Leafhoppers (including grape leafhopper and variegated leafhopper) Grape cane girdler Grape berry moth Glasswinged sharpshooter Aphids Mealybug (grape, obscure, vine) Western Grapeleaf Skeletonizer Thrips Phylloxera* (aerial form only) Banded Grape Bug* Rose Chafer* Japanese Beetle *	0.047 – 0.1	2.5 – 5.3	Begin application when treatment thresholds have been reached.  Thorough coverage is important to obtain optimum control.  For Mealybug control, apply ASSAIL 30SG Insecticide as crawlers/nymphs become active.  For Western grapeleaf skeletonizer, apply ASSAIL 30SG Insecticide as larvae are observed feeding on leaves. Apply sufficient water to provide thorough coverage of all surfaces.  For Japanese Beetle: Adult beetles will stop feeding after application and mortality will occur within a few days.  The use of spray adjuvants, such as high quality non-ionic surfactants, methylated seed or horticultural oils, may enhance coverage and plant uptake and may improve pest control. The use of stickers is not recommended. Some adjuvants can cause adverse effects, such as spotting or burn to fruit or foliage. Select an adjuvant that will be safe for the target crop. Follow adjuvant use directions. *Not for use on these pests in CA, OR, and WA

**RESTRICTIONS:** Grapes and Other Climbing Vine Small Fruits (Except Fuzzy Kiwifruit) within Crop Sub-Group 13-07F)

- Do not apply more than once every 14 days.
- Do not apply less than 3 days before harvest (PHI = 3 days)
- Do not make more than 2 applications per calendar year.
- Do not exceed a total of 0.20 lbs active ingredient (10.6 ozs product) per calendar year.

United Phosphorus, Inc.  
 630 Freedom Business Center, Suite 402  
 King of Prussia, PA 19406  
 1-800-438-6071