

---

## SUPPLEMENTAL LABELING

---

**Syngenta Crop Protection, LLC**  
P. O. Box 18300  
Greensboro, North Carolina 27419-8300

**SCP 1162A-S3 1222**

**Boundary® 6.5EC Herbicide**

**This supplemental label expires on 11/14/2025 and must not be used or distributed after this date.**

<b>Active Ingredients:</b>	<b>% by wt.</b>
S-metolachlor *: .....	58.2%
Metribuzin** .....	13.8%
Other Ingredients***: .....	28.0%
<b>Total:</b>	<b>100.0%</b>

\*CAS No. 87392-12-9

\*\*CAS No. 21087-64-9

\*\*\*Contains petroleum distillates.

Boundary® 6.5EC Herbicide is formulated as an Emulsifiable Concentrate (EC) containing 5.25 lb of S-metolachlor and 1.25 lb of metribuzin per gallon.

**KEEP OUT OF REACH OF CHILDREN**

## **WARNING/AVISO**

**EPA Reg. No. 100-1162**

**All applicable directions, restrictions and precautions on the EPA registered label are to be followed. Before using Boundary 6.5EC Herbicide as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

**syngenta**

## DIRECTIONS FOR USE

### Sugarcane (Only for use in LA, FL, and TX)

Preplant Surface, Preemergence, or Postemergence Applications

Crops (including cultivars, varieties, and/or hybrids of these)		
Sugarcane		
Application Timing	Rate (pt/A)	Use Directions
<b>Preplant Surface</b>  <b>Preemergence</b>  <b>Postemergence</b>	1.5 to 3.0 pt/A	<p>Dry weather following Boundary 6.5EC application may reduce effectiveness.</p> <p><b>Preplant Application:</b> Apply prior to planting of cane.</p> <p><b>Preemergence Application:</b> Apply after planting of cane but prior to crop emergence.</p> <p>Application can also be made after harvest of ratoon cane.</p> <p><b>Postemergence Application:</b> Spray contact with sugarcane foliage may result in minor leaf margin chlorosis and/or necrosis.</p> <p>The addition of a spray adjuvants such as crop oil concentrates (COC's) or methylated seed oils (MSO's) can increase the risk of crop injury.</p>
<b>For Weed Control:</b> Refer to <b>Section 7.1</b> for list of weeds controlled or partially controlled.		
<b>PRECAUTIONS</b>		
1) If Boundary 6.5EC Herbicide is tank-mixed with other herbicides, follow the label restrictions for the most restrictive tank-mix partner(s).		
<b>USE RESTRICTIONS</b>		
1) <b>Maximum Single Application Rate:</b> 3.0 pt/A (1.97 lb./A S-metolachlor and 0.47 lb./A metribuzin) 2) <b>DO NOT</b> apply more than two applications of Boundary 6.5EC per year. 3) <b>Minimum Application Interval:</b> 2 weeks 4) <b>Maximum Annual Rate:</b> 5.1 pt/A/year (3.34 lb./A S-metolachlor and 0.80 lb./A metribuzin) a. <b>DO NOT</b> exceed 2.0 lbs ai/A/year of metribuzin containing products in Florida. b. <b>DO NOT</b> exceed 3.0 lbs ai/A/year of metribuzin containing products in Louisiana and Texas, except Florida. c. <b>DO NOT</b> exceed 3.34 lbs ai/A/year of S-metolachlor containing products. 5) <b>DO NOT</b> use treated foliage for feed, hay, or forage. 6) <b>DO NOT</b> apply within 100 days of harvest. 7) <b>DO NOT</b> use on sand soils in Florida. 8) To assure that spray will not adversely affect adjacent sensitive nontarget plants, apply this product by aircraft at a minimum upwind distance of 400 ft. from sensitive plants.		

Boundary® and the Syngenta Logo are trademarks of a Syngenta Group Company

©2023 Syngenta