

**FIFRA Section 24(c) Special Local Need (SLN) Label**  
**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON**  
**For Control of Twospotted Spider Mite in Mint**  
**Oberon® 4 SC Insecticide/Miticide**

EPA Reg. No. 264-850

EPA SLN. No. WA-170012

**THIS LABEL FOR OBERON 4 SC INSECTICIDE/MITICIDE EXPIRES AND MUST NOT BE DISTRIBUTED OR USED IN ACCORDANCE WITH THIS SLN REGISTRATION AFTER DECEMBER 31, 2022.**

**DIRECTIONS FOR USE**

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label and this SLN label.

**MINT**

Pests Controlled	RATE PER APPLICATION fluid ounces/acre	Use Directions
Twospotted spider mite	4.0 - 8.0	Make a foliar application at the onset of infestation, prior to the establishment of a damaging population. Repeat as necessary but no closer than a 7-day interval. Application may be made by ground, by air, or by chemigation (refer to Restrictions/Precautions). Please refer to the OBERON 4 SC Section 3 label for additional information.

**RESTRICTIONS/PRECAUTIONS**

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours following application.

Pre-harvest Interval (PHI): 7 days.

Maximum OBERON 4 SC INSECTICIDE/MITICIDE allowed per 7-day interval: 8.0 fl oz per acre (0.25 lb ai/A).

Maximum OBERON 4 SC INSECTICIDE/MITICIDE allowed per crop season: 24.0 fl oz per acre (0.75 lb ai/A).

Maximum number of applications per crop season: 3.

Minimum application volume: 10.0 GPA – ground, 5.0 GPA – aerial application.

Do not apply during mint flowering. Do not apply when bees are foraging on mint.

This product may be applied through irrigation systems. Refer to the EPA registered label for chemigation directions. **Types of Irrigation Systems:** Apply OBERON 4 SC INSECTICIDE/MITICIDE only through sprinkler, including center pivot, lateral move, side roll, or overhead solid set irrigation systems. Do not apply OBERON 4 SC INSECTICIDE/MITICIDE through any other type of irrigation system.

This pesticide is toxic to fish and aquatic invertebrates. OBERON 4 SC INSECTICIDE/MITICIDE should not be used under this SLN label where impact on listed threatened or endangered species is likely. You may refer to the WSDA Natural Resources Assessment Section web site ([agr.wa.gov/PestFert/NatResources/EndangSpecies.aspx](http://agr.wa.gov/PestFert/NatResources/EndangSpecies.aspx)), or contact the Washington Department of Fish & Wildlife, National Marine Fisheries Service (NOAA Fisheries) or US Fish & Wildlife Service for information regarding aquatic species listed as threatened or endangered. Do not contaminate surface water through spray drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwater. Consult the federal label for additional restrictions and precautions to protect aquatic organisms.

### Surface Water Advisory

This product may contaminate water through runoff or drift of spray in wind. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

### WSDA Chemigation Guidance:

- Application off-site is prohibited. The chemigation application must be continuously observed whenever sensitive areas as defined in WAC 16-202-1002(44) (including but not limited to schools, parks, dwellings, occupied buildings or structures, public roadways, and waters of the state) are at risk of being exposed to drift, runoff, or overspray. In order to minimize the potential for application off-site, WSDA recommends that the product only be applied through low pressure irrigation systems (defined as 2 to 35 pounds/square inch measured at the nozzle) with a nozzle release height no higher than 3 feet above the target crop, and that end guns be disabled throughout the application.
- An inspection port or a direct access point is required, and it must be positioned immediately upstream of the irrigation mainline check valve and be of sufficient size to allow visual and manual inspection of the check valve and low pressure drain. The inspection port or access point must have a minimum diameter of four inches, unless an alternative access system is approved by WSDA (WAC 16-202-1012[1]).
- The chemigation application tank cannot be placed within 20 feet of the wellhead or other sensitive areas. Mixing or loading activities cannot occur within 20 feet of the wellhead or other sensitive areas (WAC 16-202-1008[1]).
- WSDA Chemigation Rules (WAC 16-202-1001 through WAC 16-202-1024), and information on USEPA Authorized Alternative Chemigation Safety Equipment, Distribution Uniformity and other chemigation topics are available on the WSDA web site ([agr.wa.gov/PestFert/ChemFert/default.aspx](http://agr.wa.gov/PestFert/ChemFert/default.aspx)).

**WSDA Container Disposal Guidance:** Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the WSDA Waste Pesticide Program web site ([agr.wa.gov/PestFert/Pesticides/WastePesticide.aspx](http://agr.wa.gov/PestFert/Pesticides/WastePesticide.aspx)). Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577.

For **PRODUCT USE** Information Call **1-866-99BAYER (1-866-992-2937)**.

*As with any crop-protection product, always read and follow label directions.*

*For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*

**8/8/2017**

### Section 24c SLN Registrant:

**Bayer CropScience, LP**  
**P.O. Box 12014**  
**2 T.W. Alexander Drive**  
**Research Triangle Park, NC 27709**