



# SUPPLEMENTAL LABELING

**KASUGAMYCIN    GROUP    24    FUNGICIDE**

## **KASUMIN® 2L**

**Supplemental Label to Add DIRECTIONS FOR USE on WALNUT.**

**This supplemental label expires March 1, 2021 and must not be used or distributed after this date.**

**For Agricultural Use Only**

**ACTIVE INGREDIENT:**

\*Kasugamycin Hydrochloride Hydrate ..... 2.3%

**OTHER INGREDIENTS:**..... 97.7%

**TOTAL:**..... 100.0%

\*(Equivalent to 2.0% kasugamycin)

Contains 0.168 pounds kasugamycin per gallon

**KEEP OUT OF REACH OF CHILDREN  
CAUTION / PRECAUCIÓN**

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. READ THE LABEL AFFIXED TO THE CONTAINER FOR **KASUMIN® 2L** BEFORE APPLYING. USE OF **KASUMIN 2L** ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR **KASUMIN 2L (EPA Reg. No. 66330-404)**.

**030118V002Su**

## WALNUT

Disease Control	Application Rate fl oz product (lbs ai)	Specific Use Instructions
<b>Walnut Blight</b> <i>(Xanthomonas campestris pv juglandis)</i>	<b>64 fl oz/A</b> <b>(0.084 lb ai/A)</b>	<ul style="list-style-type: none"> <li>▪ Apply at a rate of <b>64 fl oz/A in at least 100 gal of water by ground application.</b></li> <li>▪ Reduced spray volumes may be utilized for small trees where complete coverage can be obtained with a less than 100 gal of water/A.</li> <li>▪ Begin applications when conditions favor disease development.</li> <li>▪ For airblast application: To help reduce off-target drift, direct spray into the canopy, and turn off outward pointing nozzles at row ends and when spraying outer rows.</li> </ul>

### RESTRICTIONS:

- Do not apply more than 64 fl oz/A (0.084 lb ai/A) in a single application.
- Do not apply more than 256 fl oz (0.336 lb ai/A) of **KASUMIN 2L** per acre per year.
- **In California:** Do not apply more than 128 fl oz (0.168 lb ai/A) of **KASUMIN 2L** per acre per year.
- Do not make more than 4 applications of **KASUMIN 2L** per year with a minimum interval of 7 days between applications.
- **In California:** Do not make more than 2 applications of **KASUMIN 2L** per year.
- Do not apply more than 2 consecutive applications of **KASUMIN 2L** before alternating with another fungicide/bactericide of a different mode of action. If additional bactericide applications are needed, utilize an alternate product registered for this use. Two more consecutive **KASUMIN 2L** applications may be made subsequent to use of the alternate registered product.
- Do not use an alternate tree-row application method.
- Do not apply kasugamycin in orchards in which the soil has been fertilized with animal waste/manure or human biosolids.
- Animal grazing in treated areas is prohibited. The public must be notified by posting restriction signs along the perimeter of the treatment area.
- Do not apply **KASUMIN 2L** through any type of irrigation system, including chemigation.
- Do not apply **KASUMIN 2L** by aerial application.
- Preharvest Interval (PHI) = 100 days.
- Use only the specified and full-strength application rates.

### Produced For:

**ARYSTA LIFESCIENCE NORTH AMERICA, LLC**  
**15401 Weston Parkway, Suite 150**  
**Cary, NC 27513**

Arysta LifeScience and the Arysta LifeScience logo are registered trademarks of Arysta LifeScience Corporation. The **KASUMIN** logo is a registered trademark of an Arysta LifeScience Group Company.

**KASUMIN** is a registered trademark of Hokko Chemical Industry Co., Ltd.

©2018 Arysta LifeScience Group Company.