

Product Bulletin



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Intrepid Edge[®]

EPA Reg. No. 62719-666

Special 2(ee) Recommendation[†]

(For distribution and use in Alabama, Arkansas, Louisiana, Mississippi, and Tennessee only)

For Late Season Control of Soybean Looper and Other Lepidopterous pests in Soybeans

ATTENTION

[†]This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Intrepid Edge[®] insecticide before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Use of Intrepid Edge according to this Section 2(ee) recommendation is subject to all use precautions and limitations imposed by the label affixed to the container for Intrepid Edge.

Directions for Use

Refer to product label for Intrepid Edge[®] for General Use Precautions, Mixing Directions, Application Directions, and Restrictions.

Application Rate: The amount of Intrepid Edge[®] applied per acre will depend upon plant size, volume of foliage present, pest growth stage and pest infestation level. Use a lower rate in the rate range for light infestations, and a higher rate in the rate range for heavy infestations. This special recommendation is specifically for lepidopterous pests that occur when the crop is nearing maturity and less residual control is needed.

Ground Application: Apply in a minimum spray volume of 10 gpa using calibrated ground application equipment that provides thorough coverage.

Aerial Application: Apply in a minimum spray volume of 5 gpa in equipment that has been properly patterned and calibrated for environmental conditions at the site. Use higher water volumes for heavy infestations and in situations where thorough coverage is difficult to achieve.

Resistance Management: Do not make more than two consecutive applications of Group 5 or Group 18 insecticides. If additional treatments are required after two consecutive applications of Group 5 or Group 18 insecticides, rotate to another class of effective insecticides for at least one application. Consult your Dow AgroSciences representative, extension specialist, certified crop advisor, or state agricultural experiment station for information on alternative effective products to use in your area. Avoid applying Intrepid Edge[®] and other Group 5 and Group 18 insecticides to consecutive generations of all target insect pests.

Pests	Application Rate (fl oz/acre)	Application Timing
-------	----------------------------------	--------------------

beet armyworm cabbage looper fall armyworm green clover worm saltmarsh caterpillar southern armyworm soybean loopers true armyworm velvet bean caterpillar yellowstriped armyworm	3 – 4	Begin applications after the crop has reached R5 (beginning seed) when first signs of feeding damage appear or when threshold levels of feeding damage occur. Do not make applications after the crop has reached R7 (beginning maturity).
--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Restrictions:

- **Preharvest Interval:** Do not apply within 28 days of seed harvest.
- Do not apply more than 6.4 fl oz of Intrepid Edge® per acre per application.
- Do not apply more than a total of 27.75 fl oz of Intrepid Edge® (0.545 lb ai methoxyfenozide, 0.109 lb ai spinetoram) per acre per year.
- **Minimum Treatment Interval:** Do not make applications less than 4 days apart.
- **Maximum Number of Applications:** Do not make more than 4 applications per year.
- **Re-Planting Interval:** A 7-day re-planting interval is required for residues of methoxyfenozide.

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

R911-007

Issued: 08/15/16

Replaces R911-001