



## TREATED SEED

This seed has been treated with INTEGO™ Solo Fungicide which contains ethaboxam, and INOVATE® Pro Seed Protectant which contains clothianidin, ipconazole and metalaxyl at the rate recommended by the manufacturer.

## CAUTION

### DO NOT USE TREATED SEED FOR FEED, FOOD OR OIL PROCESSING

### STORE AWAY FROM FEEDS AND OTHER FOODSTUFFS

### KEEP OUT OF REACH OF CHILDREN

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Wear long-sleeved shirt, long pants and chemical-resistant gloves such as Natural Rubber  $\geq 14$  mils, Barrier Laminate, Butyl Rubber, Nitrile Rubber, Neoprene Rubber or Viton when handling treated seed.

- Excess seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agonomic practices.
- Dispose of all excess treated seed. Leftover treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements. Do not contaminate water bodies when disposing of planting equipment washwaters.
- Dispose of seed packaging in accordance with local requirements.
- Do not graze or feed soybean forage and hay to livestock.



#### HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 800-892-0099 for emergency medical treatment information.

- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading.

EPA Reg. 59639-186

059639-00186.20140603.NOTIF

INTEGO is a trademark of Valent U.S.A. Corporation

# INOVATE®

**PRO** Seed Protectant

#### FIRST AID

**If swallowed:** Call a poison control center or doctor immediately for treatment advice.

Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor.

**If in eyes:** Do not give anything to an unconscious person. Hold eye open and rinse slowly and gently with water for 15-20 minutes.

Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

Call a poison control center or doctor for treatment advice.

- Regardless of the type of application (seed treatment or foliar), do not apply more than a total of 0.2 lb (90.7 gram) clothianidin (active ingredient) per acre per season. (Based on the seed size of 3000 seeds/lb, at the label rate, the mg of clothianidin will be equal to 0.075 mg/seed).
- The compound is toxic to birds and mammals. Treated clothianidin seeds exposed on soil surface may be hazardous to birds and mammals. Cover or collect seeds spilled during loading.
- Treated seeds must be planted at a minimum depth of 1 inch.
- This compound is highly toxic to bees exposed directly (contact). Ensure that planting equipment is functioning properly in accordance with manufacturing recommendations to minimize seed coat abrasion during planting to reduce dust which can drift to blooming crops or weeds.
- This product contains metalaxyl that is known to leach through soil into groundwater under certain conditions as result of agricultural use. Use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.
- Areas planted with treated seed of soybean may be replanted immediately with bulb vegetables, canola, cereal grains, cotton, corn (field, pop and sweet), fruiting vegetables (including cucurbits), leafy vegetables (including brassica vegetables), rapeseed, root and tuber vegetables and soybean. These areas may also be replanted after 30 days with dry beans, grasses and non-grass animal feeds. These areas may also be replanted after 8 months with sugarcane. Do not plant any other crop in the treated area for at least one year after these treated seeds are planted.
- After seeds have been planted, do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. Exception: Once seeds are planted and covered by soil or other planting media, the Worker Protection Standard (WPS) allows workers to enter the treated area without restriction.
- Do not re-use bags from treated seed to handle food or feed products.
- Do not use augers used for handling treated seed to move seed for feed, food or oil processing.

EPA Reg. 59639-195

059639-001 95.20140116.FINAL

Inovate is a registered trademark of Chemtura Corporation

For **PRODUCT USE** information:

Call 1-800-6-VALENT (1-800-682-5368)

**IN CASE OF EMERGENCY ONLY:**

Call 24 Hours A Day 1-800-892-0099

**Valent U.S.A. Corporation**

P.O. Box 8025  
Walnut Creek, California 94596-8025

Form 1982-A