

# FIFRA Section 24(c) Special Local Need Label



Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Intrepid 2F

EPA Reg. No. 62719-442

EPA SLN No. MS-060014

For Distribution and Use Only in the State of Mississippi

### Control of Lepidoptera Larvae Infesting Sweet Potato

**Expiration date:** This label expires and must not be distributed or used in accordance with this SLN registration after December 31, 2028

#### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read this SLN and the label affixed to the container for this product before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of this product according to this FIFRA 24(c) Special Local Need labeling is subject to all use precautions and limitations imposed by the label affixed to the container for this product.

#### Directions for Use

Refer to product labeling for Use Precautions and Restrictions, Mixing and Application instructions.

#### Pests, Application Rates, Application Timing and Restrictions:

Pests	Application Rate (fl oz/acre)	Application Timing	Restrictions
armyworms cabbageworms loopers saltmarsh caterpillar webworms	6 - 10 (0.09 – 0.16 lb ai/acre)	Apply at egg hatch or when first signs of feeding occur. Use the higher rates for heavier infestations and under conditions in which thorough coverage is more difficult.  Under heavy infestations, continuous moth flights and/or egg masses and larvae in all stages of development, reapply to protect new growth until moth flights and/or hits subside.	<ul style="list-style-type: none"> <li>• Do not apply more than 30 fl oz of Intrepid 2F (0.48 lb ai) per acre per season.</li> <li>• Do not make more than 3 applications per acre per season.</li> <li>• <b>Pre-harvest Interval:</b> Do not harvest within 7 days of application.</li> <li>• See Rotational Crop Restrictions below.</li> </ul>

**Ground Application:** Apply a minimum of 10 gallons per acre by conventional ground equipment. Higher spray volumes may be required to provide thorough coverage to a larger, more mature crop. Equipment and spray volume should be calibrated to assure uniform coverage of infested parts of the crop.

**Aerial Application:** Apply a minimum of 10 gallons per acre. Use sufficient carrier volume to provide thorough, uniform coverage. Use of higher carrier volume is recommended for heavy infestations and in situations where thorough coverage is difficult to achieve.

**Rotational Crop Restrictions:** Crops with registered uses may be replanted at any time. All other crops grown for food or feed may be replanted after 7 days.

**Note:** Drift and runoff from applications of this product may be hazardous to sensitive aquatic invertebrates in water bodies adjacent to the treatment area. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

™®Trademarks of Corteva Agriscience and its affiliated companies

R846-112

Approved: 10/04/2023

Replaces R846-045