

CM

LIQUID CaB

A Complexed Formulation for the Correction and Prevention of Calcium & Boron Deficiencies

GUARANTEED ANALYSIS 3-0-0

Total Nitrogen (N).....	3.0%
3% Nitrate Nitrogen	
Calcium (Ca).....	5.5%
Boron (B)	0.5%
Derived from Calcium Nitrate and Boric Acid	
Wt per gal: 10.45 lb. @ 68°F Grams per liter: 1252 g @ 20°C	

GENERAL INFORMATION

CaB is designed as a foliar nutritional spray to supply Calcium and Boron in a convenient, soluble form. CaB is available and absorbed on the leaf surface or by the root system.

COMPATIBILITY: CaB is compatible with most pesticides. In mixing, always add this product to the tank last. A compatibility test is recommended if the desired combination has not been used in the past.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Follow all recommended governmental procedures for disposal of pesticide spray solutions. Rinse container, empty rinsate and spray tank. Contact county agricultural commissioner for container disposal information.

CONDITIONS OF SALE

The statements made on this label are believed to be true and accurate, but because of conditions of use which are beyond our control, the manufacturer does not make nor does it authorize any agent or representative to make any warranty or representation, expressed or implied, concerning this material or the use thereof except in conformity with the statements on the label. The buyer assumes all risk and liability resulting from improper handling, storage or use and accepts and uses this material on these conditions. If these conditions are unacceptable to the buyer, he should immediately return the unopened product to the seller.

MANUFACTURED BY:
CUSTOM AGRICULTURAL FORMULATORS
P.O. BOX 26104, FRESNO, CA 93729 800-355-4457

DIRECTIONS

For best results, spray when the crop is in an active growing state after irrigation or natural rain fall. Spray early in the morning or late afternoon. Midday sprays may not be effective because of excessive moisture evaporation.

CAUTION:

Excessive amounts of Boron may cause damage to susceptible crops. Determine plant sensitivity prior to use. Contact your local agricultural authority for information on specific crops.

Recommendations for Use: CaB may be used on the following crops: Alfalfa, Almonds, Avocados, Apples, Beets, Broccoli, Brussels Sprouts, Cabbage, Carrots, Cauliflower, Celery, Citrus, Clover, Corn, Cotton, Grapes, Lettuce, Milo, Melons, Nectarines, Rice, Pears, Peaches, Pecans, Peppers, Plums, Prunes, Potatoes, Peanuts, Sorghum, Sugar Beets, Sweet Corn, Sugarcane, Strawberries, Tobacco, Tomatoes, Turnips, Walnuts, Watermelons, Wheat and most other crops.

Vegetables:

Apply 1 pint per acre per week in the normal spray program. Increase the rate to 2 pints per acre per week prior to the time that the deficiency usually shows up.

Tree Crops:

Apply 1 to 2 pints per 100 gallons of spray. Start two to three weeks after full bloom and continue at two-week intervals. Trees must be wet thoroughly. For best results do not spray in the heat of the day.

CAUTION: This product contains boron in amounts which could be excessive for some crops. Contact your local agricultural authority for information on specific crops.

Information regarding the contents and levels of metals in this product is available by calling (800) 355-4457.

Peanuts: Apply 1 gallon per acre per application. Repeat 3 times starting at early bloom. **Cotton:** Apply a total of 1 gallon per acre with two applications of 1/2 gallon each or four treatments of 1/4 gallon per acre at 10 to 14 day intervals starting at early bloom. **Soybeans:** Apply 1/2 gallon per acre at early pod set and an additional application of 1/2 gallon per acre in 14 days.

Note: Do not exceed 1 gallon per acre per application.

Foliar Applications:

Conventional Sprayers: Use a minimum of 20 gallons of water per acre.

Low Volume Sprayers: Use 5 gallons of water per acre. If less water is used, slight burning of the foliage may occur.

Aerial Applications: Do not exceed 1 quart per gallon of water.

KEEP OUT OF REACH OF CHILDREN

WARNING



PRECAUTIONARY STATEMENTS

WARNING Hazardous to humans and animals. Causes eye and skin irritations. When mixing or applying agricultural sprays, always wear spray goggles and impervious gloves plus protective clothing as recommended on pesticide labels.

FIRST AID

If this product gets into the eyes, immediately wash the eyes with large amounts of water for at least 15 minutes, occasionally lifting the upper and lower lids. Obtain immediate medical treatment. If this product gets on the skin, immediately wash the contaminated area. Seek medical attention if skin irritation persists. If this product has been swallowed and the person is conscious, immediately give large volumes of water. Do not induce vomiting. Seek immediate medical attention.