



**FIFRA 24(c) SPECIAL LOCAL NEED LABEL (SLN)  
FOR DISTRIBUTION AND USE ONLY IN THE STATE OF CALIFORNIA**

**For Aerial Application to Almonds; Citrus Fruit Crop Group 10-10; Stone Fruit,  
Crop Group 12-12 with Standing Water**

**Quadris Top® Fungicide**

**EPA Reg. No. 100-1313**

**EPA SLN No. CA-210001**

**This label expires and must not be distributed or used in accordance with this SLN  
registration after December 31, 2025.**

**DIRECTIONS FOR USE**

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- This state-specific Section 24(c) labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, and precautions on the EPA-registered Quadris Top, EPA Reg No. 100-1313 label and this label.

Crop	Target Diseases	Use Rate fl oz product/A	Remarks
<b>Almonds</b>	Alternaria Leaf Spot ( <i>A. alternata</i> ) Anthracnose ( <i>Colletotrichum acutatum</i> ) Blossom Blight ( <i>Monilinia</i> spp.) Leaf Blight ( <i>Seimatosporium lichenicola</i> ) Leaf Rust ( <i>Tranzschelia discolor</i> ) Scab ( <i>Venturia carpophila</i> ) Shot Hole ( <i>Wilsonomyces carpophilus</i> )	12-14	For blossom blight, begin applications at early bloom and continue through petal fall. Make no more than 2 sequential applications before alternating to another fungicide with a different mode of action. For all other diseases, begin applications prior to disease onset when conditions are conducive for disease. Apply Quadris Top Fungicide on a 14- to 21-day schedule making no more than 2 sequential applications before alternating to another fungicide with a non-QoI (Group 11) mode of action. If monitoring or history indicates the presence of Alternaria, apply 14 fl oz/A of Quadris Top Fungicide in the late spring (mid-April to beginning of May) and then repeat the treatment 2-3 weeks later. The addition of a spreading/penetrating type adjuvant such as a non-ionic based surfactant or crop oil concentrate or blend is recommended. If disease pressure is high, use the shortest interval and highest rate.

**Application:** For best results, sufficient water volume must be used to provide thorough coverage. Quadris Top Fungicide can be only applied by aerial application. For aerial applications, use a minimum of 10 gal/A of water.

**Specific Use Restrictions:**

**1)** Do not apply more than 56 fl oz/A/year of Quadris Top Fungicide (0.73 lb azoxystrobin and 0.46 lb difenoconazole). **2)** Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products. **3)** Do not apply more than 1.5 lb ai/A/year of azoxystrobin-containing products. **4)** Do not apply within 28 days of harvest (28-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A	Remarks
<b>Citrus Fruit Crop Group 10-10</b>  Grapefruit Lemon Lime Orange (Sour and Sweet) Tangerine  Including all cultivars and/or hybrids of these  See complete list of citrus fruit crops below.	Greasy Spot (Mycosphaerella citri)	10-15.4	Quadris Top Fungicide applications should begin prior to disease development and continue throughout the season on 7- to 21- day intervals following the resistance management guidelines. Applications may be made by ground or air. An adjuvant may be added at recommended rates. A horticultural spray oil should be used to improve control of greasy spot. The addition of a spreading/ penetrating type adjuvant such as a non-ionic based surfactant or crop oil concentrate or blend is recommended. If disease pressure is high, use the shortest interval and highest rate. Make no more than 2 sequential applications before alternating to another fungicide with a non-QoI (Group 11) different mode of action. Do not make more than 4 applications of Quadris Top Fungicide or other Group 11 fungicides per season.
	Alternaria Leaf and Fruit Spot ( <i>Alternaria citri</i> ) Anthracnose ( <i>Colletotrichum</i> spp.) Black Spot ( <i>Guignardia citricarpa</i> ) Greasy Spot Rind Blotch ( <i>Mycosphaerella citri</i> ) Melanose ( <i>Diaporthe citri</i> ) Phomopsis Stem-End Rot ( <i>Phomopsis citrii</i> ) Post-Bloom Fruit Drop (PFD) ( <i>Colletotrichum acutatum</i> ) Scab ( <i>Elsinoe fawcettii</i> )	15.4	Quadris Top Fungicide applications should begin prior to disease development and continue throughout the season on 7- to 21- day intervals following the resistance management guidelines. Applications may be made by ground or air. An adjuvant may be added at recommended rates. A horticultural spray oil should be used to improve control of greasy spot. If disease pressure is high, use the shortest interval. Make no more than 2 sequential applications before alternating to another fungicide with a non-QoI (Group 11) different mode of action. Do not make more than 4 applications of Quadris Top Fungicide or other Group 11 fungicides per season. The addition of a spreading/penetrating type adjuvant such as a non-ionic based surfactant or crop oil concentrate or blend is recommended.

**Application:** For best results, sufficient water volume must be used to provide thorough coverage. Quadris Top Fungicide can be only applied by aerial application. For aerial applications, use a minimum of 10 gal/A of water.

**Complete List of Citrus Fruit Crops:** Australian desert lime; Australian finger lime; Australian round lime; Brown River finger lime; Calamondin; Citron; Citrus hybrids (*Citrus* spp., *Eremocitrus* spp., *Fortunella* spp., *Microcitrus* spp., and *Poncirus* spp.); Grapefruit; Japanese summer grapefruit; Kumquat; Lemon; Lime; Mediterranean mandarin; Mount White lime; New Guinea wild lime; Orange, sour; Orange, sweet; Pummelo; Russell River lime; Satsuma mandarin; Sweet lime; Tachibana orange; Tahiti lime; Tangelo; Tangerine (Mandarin); Tangor; Trifoliate orange; UniQ fruit; cultivars, varieties and/or hybrids of these.

**Specific Use Restrictions:**

**1)** Do not use Quadris Top Fungicide in citrus plant propagation nurseries. **2)** Do not apply more than 61.5 fl oz/A/year of Quadris Top Fungicide (0.80 lb azoxystrobin and 0.50 lb difenoconazole). **3)** Do not apply more than 0.5 lb ai/A/year of difenoconazole-containing products. **4)** Do not apply more than 1.5 lb ai/A/year of azoxystrobin-containing products. **5)** May be applied the day of harvest (0-day PHI).

Crop	Target Diseases	Use Rate fl oz product/A	Remarks
<b>Stone Fruit, Crop Group 12-12</b> Apricots Cherries, Sweet Cherries, Tart Nectarines Peaches Plums Plumcot Prunes  Including all cultivars and/or hybrids of these	Alternaria Spot and Fruit Rot <i>(A. alternata)</i>  Anthracnose <i>(Colletotrichum spp.)</i>  Brown Rot Blossom Blight and Fruit Rot <i>(Monilinia fructicola, M. laxa)</i>  Leaf Rust <i>(Tranzschelia discolor)</i>  Powdery Mildew <i>(Sphaerotheca pannosa, Podosphaera clandestina)</i>  Scab <i>(Cladosporium carpophilum)</i>  Shot Hole <i>(Wilsonomyces carpophilus)</i>	12-14	For brown rot blossom blight, begin applications at early bloom and continue through petal fall. For brown rot on fruit, apply as needed a maximum of two sprays during the pre-harvest period up to the day of harvest (minimum of a 7-day retreatment interval). If high inoculum and severe disease conditions persist, apply a registered fungicide that is non-Group 11 or non-Group 9. For all other diseases, follow the brown rot blossom blight schedule. Make additional applications on a 10- to 14-day interval from the end of petal fall to harvest. The addition of a spreading/penetrating type adjuvant such as a non-ionic based surfactant or crop oil concentrate or blend is recommended. If disease pressure is high, use the shortest interval and highest rate.

**Application:** For best results, sufficient water volume must be used to provide thorough coverage. Quadris Top Fungicide can be only applied by aerial application, For aerial applications use a minimum of 10 gal/A of water.

**Complete List of Stone Fruit Crops:** Apricot; apricot, Japanese; capulin; cherry, black; cherry, Nanking; cherry, sweet; cherry, tart; Jujube, Chinese; nectarine; peach; plum; plum, American; plum, beach; plum, Canada; plum, cherry; plum, Chickasaw; plum, Damson; plum, Japanese; plum, Klamath; plum, prune; plumcot; sloe; cultivars, varieties, and/or hybrids of these.

**Specific Use Restrictions:**

**1)** Do not apply more than 56 fl oz/A/year of Quadris Top Fungicide (0.73 lb azoxystrobin and 0.46 lb difenoconazole). **2)** Do not apply more than 0.46 lb ai/A/year of difenoconazole-containing products. **3)** Do not apply more than 1.5 lb ai/A/year of azoxystrobin-containing products. **4)** Quadris Top Fungicide may be applied on the day of harvest (0-day PHI).

**Requirements:**

1. For applications to orchards that have standing water due to a storm event.
2. Water must be standing and will not flow into state waterways where fish and wildlife are likely to be affected.
3. Applications are made in accordance with all other label directions.
4. The County Agriculture Commissioner must assure the required conditions are met.

**Restrictions:**

1. No fungicide application shall occur if:
    - a. Soil moisture is at field capacity and a storm event, forecasted by the National Oceanic and Atmospheric Administration (NOAA) or National Weather Service (NWS), is to occur within 48 hours following the application; or
    - b. A storm event likely to produce runoff from the treated area is forecasted by NOAA/NWS to occur within 48 hours following the application.
  2. Applications are not allowed if water is flowing off the intended application site.
  3. Standing water will not be pumped or removed from the orchard after the fungicide application.
  4. Do not apply this product through any type of irrigation system.
- 

Quadris Top® registered trademark of a Syngenta Group Company

©2019 Syngenta

24(c) Registrant:

Syngenta Crop Protection, LLC.

P. O. Box 18300

Greensboro, North Carolina 27419

Telephone: 1-800-334-9491

Label Code: CA1313191AA0819