



Bayer CropScience LP  
P.O. Box 12014  
2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709  
1-866-99BAYER (1-866-992-2937)  
<http://www.bayercropscienceus.com>

## ALIETTE® WDG brand Fungicide

For use on Tomatoes for suppression of  
Bacterial Spot

EPA Reg. No. 264-516

# PRODUCT BULLETIN

## For Use and Distribution Only in the State of Florida

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA. It is a violation of Federal and State Law to use this product in a manner inconsistent with its label. This product bulletin must be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

ALIETTE® WDG Fungicide applied in conjunction with good cultural management practices will control damping off caused by *Pythium spp.*, root rots caused by *Phytophthora spp.* and will suppress bacterial spot (*Xanthomonas campestris*).

DISEASE	APPLICATION METHOD	RATE lbs product	APPLICATION PROGRAM
Root rot ( <i>Phytophthora spp.</i> ) Damping off ( <i>Pythium spp.</i> ) Bacterial spot ( <i>Xanthomonas campestris</i> ) ( <b>Suppression only</b> )	Foliar	2.5-5.0/Acre	Begin applications at the 2-4 leaf stage of growth for direct seeded tomatoes or immediately after transplanting to the field. Continue applications on a 7-14 day interval if conditions are favorable for disease development. Apply ALIETTE® WDG in a sufficient volume of water to wet the entire plant canopy, but do not apply in less than 10 gallons of water per acre.  ALIETTE® WDG should not be applied within 14 days of harvest (PHI).  Do not apply more than 20 lbs of product per acre per season.  With moderate disease pressure the higher labeled rate of ALIETTE® WDG and a shorter spray interval should be used (7 days).

**Note:** Bayer CropScience has determined that phytotoxicity from solubilized copper may occur if products containing copper are tank-mixed with ALIETTE® WDG or if unbuffered ALIETTE® WDG is applied to foliage with copper residues.

**Do not tank-mix with copper fungicides. When applied prior to or after copper fungicides, the pH of ALIETTE® WDG should be raised to 6.0 or above with the addition of an alkaline buffer such as Potassium Carbonate (3 lbs Potassium Carbonate to 5 lbs ALIETTE® WDG) or DiAmmonium Phosphate (5 lbs DiAmmonium Phosphate to 5 lbs ALIETTE® WDG).**

ALIETTE is a registered trademark of Bayer.