



Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)
<http://www.bayercropscienceus.com>

ALIETTE® WDG brand Fungicide

Foliar spray to apples for the suppression
of *Alternaria* Leaf Spot

EPA Reg. No. 264-516

PRODUCT BULLETIN

For Use and Distribution Only in the States of Georgia, North Carolina, South Carolina, and Virginia

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA. It is a violation of Federal and State Law to use this product in a manner inconsistent with its label. This product bulletin must be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

ALIETTE® WDG applied as a foliar spray in conjunction with good cultural management practices will suppress *Alternaria* Leaf Spot (*Alternaria mali*) on apple.

DISEASE	RATE LBS. PRODUCT	APPLICATION METHOD
<i>Alternaria</i> Leaf Spot (<i>Alternaria mali</i>)	2.5 – 5.0/Acre	Begin foliar applications prior to disease development (normally mid-May). An additional 2 or 3 applications should be made at 14 day intervals. Sufficient spray volume for thorough coverage of foliage and fruit is essential.

RESTRICTIONS AND LIMITATIONS:

Do not apply within 14 days of harvest (PHI = 14 days).

Do not apply within 2-3 weeks of leaf senescence.

Do not exceed 500 GPA.

Application Note: Do not apply more than 5.0 lbs of ALIETTE® WDG per acre per application. Do not exceed 20 lbs per acre per season. Applications for Fire Blight and Blister Spot control should occur early in the season. Applications for *Phytophthora* spp. control should be delayed until 30 days after last Fire Blight or Blister Spot application.

Do not graze livestock on floor of treated orchards.

Note: Bayer CropScience has determined that phytotoxicity from solubilized copper may occur if products containing copper are tank-mixed with ALIETTE® WDG or if unbuffered ALIETTE® WDG is applied to foliage with copper residues.

Do not tank-mix with copper compounds. When applied prior to or after copper compounds, the pH of ALIETTE® WDG should be raised to 6.0 or above with the addition of an alkaline buffer such as Potassium Carbonate (3 lbs Potassium Carbonate to 5 lbs ALIETTE® WDG) or DiAmmonium Phosphate (5 lbs DiAmmonium Phosphate to 5 lbs ALIETTE® WDG).

Adjuvants which enhance pesticide penetration should not be used with ALIETTE® WDG.

ALIETTE is a registered trademark of Bayer.