CROP MAX - N 15-0-2

15-0-2

GUARANTEED ANALYSIS

Total Nitrogen (N)..... 4.6% Ammoniacal Nitrogen

> 3.5% Nitrate Nitrogen 6.9% Urea Nitrogen

Soluble Potash (K20)......2.00% Iron(Fe)..... .05% Chelated Iron

Zinc(Zn).....

2.50% Chelated Zinc

Derived from urea ammonium nitrate, anhydrous ammonia.potassium nitrate, boric acid, Iron Ethydiaminetetraacetate(Fe EDTA), and Zinc Ethydiaminetetraacetate(Zn EDTA).

Chlorine (CI), Maximum.....

Weight per gal: 10.5 lbs/gal at 68*f

GENERAL INFORMATION

CROP MAX-N is formulated for field, vegetable, ro, fruit,

CROP MAX-N may be applied alone or with fertilizers or pesticides. It may be applied either to the plant foliage or

CROP MAX-N is compatible with most pesticides and fertilizers, except lime-sulfur, basic metal sulfates, metal recommended if the desired combination has not been used in the past.

Shake or stir contents before use. Add to spray tank last.

Information regarding the content and level of metals in this product is available by calling (800) 355-4457.

SPECIMEN LABEL

Harmful if swallowed. May cause eye and skin irritation. In case of contact, immediately flush eyes or skin with plenty of

CAUTION: This product contains boron in amounts which could be excessive for some crops. Contact your local agricultural authority for information.

CONDITIONS OF SALE

Seller warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. Weather, crop conditions, timing and methods of applications, mixtures with other chemicals not specifically recommended in the use of this product are beyond the control of the saller.

RATES FOR USE

All crops:

Maintenance: 1-2 gt/acre Mild deficiency: 2-4 qt/acre
High deficiency: 4 qt/acre
Field, row, vegetable crops: Use the lower rates
indicated above.
Fruit, tree, vine crops: Use the higher rates
indicated above.

CAUTION: Do not apply when crops are under stress. DO NOT apply to apples, apricots, nectarines, plums, prunes and other smooth-skinned fruit or grapes during bloom. Do not use on apples during drying conditions. NOTE: When rates of over 2 quarts are called for, split or multiple applications are preferred. **Drip Irrigation Application: Apply the suggested** rate of this product in multiple applications 10-14 days apart throughout the growing season of the crop.

Aerial Application: Apply 2 quarts per 8-10 gallons of water per acre.

MANUFACTURED BY: CUSTOM AGRICULTURAL FORMULATORS, P.O. BOX 26104, FRESNO, CA. 93729