

SUPPLEMENTAL LABEL

Winfield Solutions, LLC
P.O. Box 64589
St. Paul, MN 55164-0589

GROUP 9 HERBICIDE

CORNERSTONE® 5 Plus
HERBICIDE

EPA Reg. No. 1381-241

FOR USE ON ROUNDUP READY FLEX COTTON

This supplemental label expires on 5/30/2019 and must not be used or distributed after this date.

ACTIVE INGREDIENT:

*Glyphosate, N-(phosphonomethyl) glycine, in the form of its isopropylamine salt 53.8%

OTHER INGREDIENTS: 46.2%

TOTAL 100.0%

* Contains 660 grams per liter or 5.5 pounds per U.S. gallon of the active ingredient glyphosate, in the form of its isopropylamine salt. Equivalent to 480 grams per liter or 4 pounds per U.S. gallon of the acid, glyphosate.

KEEP OUT OF REACH OF CHILDREN
CAUTION

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Before using Cornerstone 5 Plus according to this Supplemental Label, read and follow all applicable directions, restrictions, precautions and conditions of sale and warranty information on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling must be in the possession of the user at the time of pesticide application.

The directions for use of this product provided in this Supplemental Label are specific to and may only be used with varieties designated as Roundup Ready Flex cotton. The use of over-the-top applications described in this Supplemental Label on cotton varieties other than Roundup Ready Flex cotton will cause crop injury and reduced yields. **DO NOT** combine the directions for use in this Supplemental Label with those in the "Cotton with the Roundup Ready Gene" section of the container label, or with any other Roundup Ready cotton or Roundup Ready Flex cotton directions for use on labeling for this or other glyphosate-containing products. Drift of this product from an application made to Roundup Ready Flex cotton onto adjacent fields of post 4-leaf (node) Roundup Ready cotton may cause extensive crop injury, including boll loss, delayed maturity and/or yield loss.

TYPES OF APPLICATION: Preplant, At-Planting, Pre-emergence, Postemergence (In-crop), Preharvest

Maximum Application Rates	
Combined total per year for all applications	6 quarts per acre
Total of all Preplant, At-Planting, Pre-emergence applications	3.75 quarts per acre
Total of all in-crop applications from cracking to 60 percent open bolls	4.5 quarts per acre
Total of all in-crop applications between layby and 60 percent open bolls	1.5 quarts per acre
Total of all in-crop applications from 60 percent open bolls to 7 days prior to harvest	1.5 quarts per acre
Total of all in-crop applications from emergence through harvest	4.5 quarts per acre

When applied in accordance with this Supplemental Label Cornerstone 5 Plus will control labeled annual grasses and broadleaf weeds in Roundup Ready Flex cotton. Many perennial weeds will be controlled or suppressed with one or more applications of this product. To maximize yield potential, eliminate competing weeds early. For specific rates of application and additional instructions for control of various annual and perennial weeds, refer to the "ANNUAL WEEDS RATE TABLE" and the "PERENNIAL WEEDS RATE TABLES" sections of the container label.

See the "ROUNDUP READY CROPS" section of the container label for additional information regarding the use of this product in Roundup Ready crops.

The maximum use rates stated throughout this Supplemental Label apply to this product combined with the use of all other herbicides containing glyphosate or sulfosate as the active ingredient, whether applied as mixtures or separately. Calculate the application rates and ensure that the total use of this and other glyphosate or sulfosate containing products does not exceed stated maximum use rates.

Preplant, At-Planting, Pre-emergence

USE INSTRUCTIONS: This product may be applied before, during or after planting Roundup Ready Flex cotton.

RESTRICTIONS: The maximum quantity of this product that may be applied for all preplant, at-planting and pre-emergence applications combined is 3.75 quarts per acre per season.

Postemergence (In-crop)

USE INSTRUCTIONS: Use an initial application rate of 24 fluid ounces (0.75 quarts) per acre to control or suppress 1 to 3 inch tall annual grasses and broadleaf weeds. This product may be applied postemergence to Roundup Ready Flex cotton using ground application equipment at rates up to 36 fluid ounces (1.125 quarts) per acre per application. In addition to broadcast application, post-directed spray equipment may be used to achieve more thorough weed coverage.

PRECAUTION: In-crop application rates above 24 fluid ounces (0.75 quarts) per acre made alone or with the addition of other crop chemical products containing surfactant may cause a crop response including leaf speckling or leaf necrosis.

DO NOT EXCEED A SURFACTANT CONCENTRATION 0.5% BY WEIGHT (2 QUARTS PER 100 GALLONS OF SPRAY SOLUTION) WHEN MAKING AN OVER-THE-TOP IN-CROP APPLICATION TO ROUNDUP READY FLEX COTTON.

RESTRICTIONS: The maximum single, in-crop application rate of this product to Roundup Ready Flex cotton using ground application equipment is 36 fluid ounces (1.125 quarts) per acre. Do not exceed a maximum rate of 24 fluid ounces (0.75 quarts) of this product per acre when using aerial application equipment. Between layby and 60 percent open bolls, the maximum combined total application rate of this product is 48 fluid ounces (1.5 quarts) per acre. The maximum combined total of all applications of this product made from crop emergence to 60 percent open bolls must not exceed 4.5 quarts per acre.

DO NOT ADD ADDITIONAL SURFACTANT OR ADDITIVES CONTAINING SURFACTANT TO THIS PRODUCT FOR OVER-THE-TOP APPLICATION TO ROUNDUP READY FLEX COTTON.

Preharvest

USE INSTRUCTIONS: Up to 48 fluid ounces (1.5 quarts) of this product per acre may be applied to Roundup Ready Flex cotton for annual and perennial weed control prior to harvest after 60 percent boll crack.

NOTE: This product will not enhance the performance of harvest aids when applied to Roundup Ready Flex cotton.

PRECAUTION: DO NOT EXCEED A SURFACTANT CONCENTRATION 0.5% BY WEIGHT (2 QUARTS PER 100 GALLONS OF SPRAY SOLUTION) WHEN MAKING AN OVER-THE-TOP IN-CROP APPLICATION TO ROUNDUP READY FLEX COTTON.

RESTRICTIONS: Allow a minimum of 7 days between application and harvest of Roundup Ready Flex cotton.

ATTENTION: USE OF THIS PRODUCT IN ACCORDANCE WITH LABEL DIRECTIONS IS EXPECTED TO RESULT IN NORMAL GROWTH OF ROUNDUP READY FLEX COTTON. HOWEVER, DUE TO THE SENSITIVITY OF COTTON FRUITING TO VARIOUS ENVIRONMENTAL CONDITIONS, AGRONOMIC PRACTICES AND OTHER FACTORS, IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS ASSOCIATED WITH THIS PRODUCT, EVEN WHEN APPLICATIONS ARE MADE IN CONFORMANCE WITH THE LABEL SPECIFICATIONS. IN SOME CASES, THESE FACTORS CAN RESULT IN BOLL LOSS, DELAYED MATURITY AND/OR YIELD LOSS.