



P.O. Box 12014 ♦ 2 T.W. Alexander Drive ♦ Research Triangle Park ♦ North Carolina 27709

FIFRA Section 24(c) Special Local Need (SLN)

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF OREGON

Proline 480 SC Fungicide

EPA Reg. No. 264-825

EPA SLN. No. OR-150016

For Suppression of Fusarium Wilt (*Fusarium* spp.) in Spinach Grown For Seed

THIS LABEL IS VALID UNTIL DECEMBER 31, 2020 OR UNTIL OTHERWISE AMENDED, WITHDRAWN, CANCELED, OR SUSPENDED.

ACTIVE INGREDIENT:

Prothioconazole: 2-[2-(1-Chlorocyclopropyl)-3-(2-chlorophenyl)-2-hydroxypropyl]-1,2-dihydro-3H-1,2,4-triazole-3-thione	41.00%
---	--------

OTHER INGREDIENTS:59.00%

TOTAL:

100.00%

KEEP OUT OF REACH OF CHILDREN
CAUTION

ENVIRONMENTAL HAZARDS

This product is toxic to estuarine/marine invertebrates, and freshwater/estuaries/marine aquatic plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Prothioconazole-desthio (a degradate of prothioconazole) is known to leach through soil into ground water under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

Surface Water Advisory

Drift and runoff are hazardous to aquatic organisms in water adjacent to treated areas. This product has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. This label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label for Proline 480 SC Fungicide (EPA Reg. No. 264-825) and this SLN label.

Spinach Grown For Seed Production	
PEST SUPPRESSION	RATE PER APPLICATION fluid ounces/acre
Fusarium Wilt (<i>Fusarium</i> spp.)	5.7
RESTRICTIONS Do not apply to spinach grown for food or feed, nor allow plant components to be utilized for human food or animal feed consumption. Do not apply within 14 days of harvest Maximum Proline 480 SC Fungicide allowed per year: 17.1 fl oz per acre Apply in a concentrated band at planting and repeat band applications as needed with a minimum 21-day interval between applications.	
NOTES Use in rotation in resistance management programs where additional products are available Please refer to the Proline 480 SC Fungicide Section 3 label for additional information	

Special Crop Use Restrictions

The pesticide applicator, the producer of the crop, and the seed conditioner must be aware that use of this product according to this labeling is deemed a non-feed/non-food use by the Oregon Department of Agriculture, and is regulated by the Oregon Administrative Rule (OAR) 603-057-0535, Pesticide Use On Crops Grown For Seed. If the applicator of this pesticide is not the producer, the applicator must provide a copy of this labeling to the producer of the crop. Producers of this crop who use this product, or cause the product to be used on a field they operate, must provide a copy of this pesticide label to the seed conditioner.

This pesticide does not have an established pesticide residue tolerance for this crop. Consequently, no portion of this seed may be used or distributed for food or feed for 1 year (365 days) after the last application of this product. This restriction pertains to, but is not limited to, green chop, forage, hay, pellets, meal, whole seed, cracked seed, straw, roots, bulbs, foliage or seed screenings, and to the grazing of the crop field, stubble or regrowth. All seed screenings shall be disposed of in such a manner that screenings cannot be distributed or used for food or feed purposes, as indicated in OAR 603-057-0535. Additional regulations concerning seed screenings are stated in OAR 603-057-0535.

Any seed from a field treated with this pesticide product shall bear specific and conspicuous container labeling, or if shipped in bulk, on the shipment invoice or bill of lading. The labeling shall contain the following statement:

"This seed was produced using one or more products for which the United States Environmental Protection Agency has not established pesticide residue tolerances. This seed, in whole, as sprouts, or in any form may not be used for human consumption or animal feed. Failure to comply with this condition may violate requirements of the Federal Food and Drug Administration, the Oregon Department of Agriculture and other regulatory agencies."

IMPORTANT—READ BEFORE USE

PROLINE 480 SC FUNGICIDE, WHEN USED ON SPINACH GROWN FOR SEED UNDER THIS SECTION 24(C) LABEL HAS NOT BEEN TESTED UNDER ALL CONDITIONS AND IN ALL TANKMIX SCENARIOS. BAYER RECOMMENDS THAT THE USER AND/OR GROWER TEST PROLINE 480 SC FUNGICIDE IN ORDER TO DETERMINE ITS COMPATIBILITY WITH THE INTENDED USE. **BECAUSE OF THE RISK OF CROP INJURY, CROP YIELD REDUCTION, AND/OR CROP LOSS, ALL SUCH USE IS AT USER'S AND/OR GROWER'S RISK.**

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).