

# REGALKADE® 50

## Weeds and Feeds in One Application

---

### ACTIVE INGREDIENTS:

Prodiamine [2,4-dinitro-N3,N3-dipropyl-6-(trifluoromethyl)-1,3-benzenediamine] ..... 0.5%

OTHER INGREDIENTS: ..... 99.5%

TOTAL ..... 100.00%

---

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

---

RegalKade is a pre-emergent herbicide (prodiamine) coated on a dry fertilizer carrier for control of crabgrass, annual bluegrass, foxtail, common chickweed, henbit, knotweed and prostrate spurge, and many other annual grass and broadleaf weeds in established turfgrasses, lawns and landscape ornamentals.

---

EPA Reg. No. 48234-5

EPA Est No. 48234-GA-I



Regal Chemical Company // Alpharetta, GA 30004

FIRST AID	
<b>IF ON SKIN:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-621-5208 for emergency medical treatment information.	

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS
Causes moderate eye irritation. Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.
PERSONAL PROTECTIVE EQUIPMENT (PPE)
Applicators and other handlers must wear: <ul style="list-style-type: none"> <li>- Long-sleeved shirt and long pants</li> <li>- Waterproof gloves</li> <li>- Shoes plus socks</li> </ul> Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS:
Users should: <ul style="list-style-type: none"> <li>• Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet</li> <li>• Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.</li> <li>• Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.</li> </ul>

#### ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limit of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

If material is spilled, contain and sweep up material of spill for disposal or use per label instructions. Wash cleanup equipment to prevent contamination elsewhere. Excess quantities above normal application rates could result in damage to plants subject to spill. Water-in remainder, disc under, cover with layer of soil, or leave exposed to sunlight, as deemed appropriate to the situation.

#### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, whether directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**NOT** for use on plants being grown (1) for sale or other commercial use, (2) for commercial seed production, or (3) for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

**DO NOT** graze or feed livestock forage cut from areas treated with this product.

**DO NOT** apply this product through any type of irrigation system.

**DO NOT** apply aerially.

**DO NOT** apply to golf course putting greens.

#### GENERAL INFORMATION

RegalKade 50 is a pre-emergent herbicide on a fertilizer carrier for dryadvance of weed seed germination. RegalKade 50 controls susceptible weeds by inhibiting growth of roots and shoots as the weed seeds germinate in turf and landscapes.

RegalKade 50 is intended for use on established turfgrasses to control many annual weeds. The maximum use rates vary according to specific turf species, and the optimum use rates depend on the weed species to be controlled. For most effective weed control RegalKade 50 must be incorporated by 1/2 inch of rainfall, irrigation or shallow mechanical incorporation. Incorporation should take place as soon as possible after application and should not be delayed past 14 days from the date of application.


#### APPLICATION DIRECTIONS

Apply RegalKade 50 at recommended rates with granular spreader equipment. Be sure spreader is properly calibrated and suitable for uniform application.

#### ESTABLISHED TURF

RegalKade 50 is a pre-emergence herbicide that, when properly applied, will control grass and broadleaf weeds in established turf as listed on this label. The species of turfgrass determines the maximum amount of RegalKade 50 that may be applied during the year. Most effective weed control in turfgrasses will be obtained when RegalKade 50 is activated by at least 1/2 inch of rainfall or irrigation within 14 days of application and prior to weed germination, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days following application. See map below for approximate crabgrass germination dates.

#### CRABGRASS SEED GERMINATION DATES



#### USE PRECAUTIONS

The following use precautions apply to the use of this product in turfgrasses and lawns:

- Application of RegalKade 50 may thin emerged annual bluegrass and newly over-seeded grasses.
- **DO NOT** apply to OVERSEEDDED TURF for 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedling is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.
- **DO NOT** apply to newly set sod until the following year.
- Application of RegalKade 50 to turf stressed by drought, low fertility, or pest damage may result in turf injury. Disturbing the herbicide barrier with cultural practices, such as disking, may result in reduced weed control.
- **DO NOT** apply this product to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.

#### MAXIMUM ANNUAL TURFGRASS RATES

RegalKade 50 must not be applied above a maximum annual rate of 300 lbs product/A (1.5 lbs. a.i./A).

#### TURF SPECIES AND RATES

For season long control RegalKade 50 may be applied as a single application, or if needed followed by a second application for increased length of control. The length of time of residual weed control provided by this product is related to the rate applied. Use the higher rate within the listed range for longer control. Longer control is desirable in areas where crabgrass germinates between January 1 and April 20 (See Crabgrass Seed Germination Dates Map).

Select maximum use rate based on turfgrass species. All applications must be made prior to germination of the target weed seeds. The product will not control established weeds.

**MAXIMUM RATE/CALENDAR YEAR of RegalKade 50 by Turfgrass Species<sup>1</sup> Table**

Turf Species:	Lbs. RegalKade 50/A	Lbs. ai/A
Bermudagrass	300	1.5
Bahiagrass		
Centipedegrass		
Seashore Paspalum		
Tall Fescue (including turf-type)		
Zoysia	200	1.0
Buffalograss		
Kentucky Bluegrass		
Perennial Ryegrass		
Creeping Red Fescue	150	0.75
St. Augustinegrass		
Creeping Bentgrass	132	0.65

<sup>1</sup>These are the maximum rate per calendar year by species limitations.

- **Do not** apply more than 300 lbs. RegalKade 50 per calendar year.
- Use higher rates of RegalKade 50 to achieve higher levels of fertility and longer periods of weed control for each turf type, but do not exceed the maximum application rates specified in the MAXIMUM RATE/CALENDAR YEAR of RegalKade 50 by Turfgrass Species Table.

**WEEDS CONTROLLED**

When used at the recommended rates (see MAXIMUM RATE/CALENDAR YEAR of RegalKade 50 by Turfgrass Species Table) this product will control:

Barnyardgrass	Knotweed
Bluegrass, Annual (Poa annua)	Lambsquarter, Common
Carpetweed	Pigweed
Chickweed, Common	Purslane, Common
Crabgrass (large, smooth)	Rescuegrass <sup>3</sup>
Foxtails, Annual	Shepherd's Purse <sup>2</sup>
Goosegrass <sup>1</sup>	Signalgrass, Broadleaf
Henbit	Spurge, Prostrate
Johnsongrass (from seed)	Woodsorrel, Yellow (from seed)

<sup>1</sup>In many areas a single application of 132 to 300 lb/A of this product will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, apply an initial treatment of 132 to 200 lbs/A followed after 60 to 90 days by a second application. Do not

**TOLERANT ORNAMENTAL SPECIES**

RegalKade 50 will not harm most established ornamental trees, shrubs and flowers. The species listed below are tolerant to this product. Best results will be obtained when this product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort

Scientific Name	Common Name
Abies spp.	Fir Species** (balsam, Fraser, Nobel, etc.)
Abelia grandiflora	Abelia: Sherwood
Acer palmatum	Japanese Maple
A. platanoides	Norway Maple
Acilliea sp.	Yarrow: King Edward
Actinidia chinensis	Kiwi*
Agapanthus africanus	Lily-of-the-Nile (African Lily)
Agapanthus Orientalis	
Akebia quintata	Five-Leaf or Chocolate Vine
Allium cernuum	Lady's Leek, Nodding Onion
Anemone hybrida	Japanese Anemone
Aquilegia sp.	Aquilegia: Red and Gold
Arctostaphylos densiflora	Vine Hill Manzanita
Arctotheca calendula	Cape Weed
Artemesia sp.	Wormwood; Silver Mound, Castle
Aster sp.	Aster: Bonny Blue, Purple Dome
Aster X Frikartii	

Scientific Name	Common Name
Athrium Filiz-femina	Lady Fern; Fern Lady
Scientific Name	Common Name
Aucuba japonica	Japanese Aucuba
Begonia sp.	Fibrous Begonia: Hardy Grandis
Berberis gladwynensis	Barberry
B. Julianae	Wintergreen Barberry
B. mentorensis	Mentor Barberry
B. Thunbergii	Japanese Barberry
B. verruculosa	Warty Barberry
Bergenia cordifolia	
Boltonia asteroides	Snowbank
Bougainvillea sp.	Bougainvillea
Buddleia davidii	Butterfly-Bush (Dwarf Blue); Royal Red
Buxus microphylla	Japanese Boxwood
Callistemon viminalis	Weeping Bottlebrush
Calluna vulgaris	Scotch Heather
Campanula carpatica	Tussock Bellflower, (White Clips)

Scientific Name	Common Name
Campis X Tagliabuana	Trumpet Creeper; Trumpet Flower; Madame Galen
Carpobrotus edulis	Hottentrot Fig (Ice Plant)
Cassia artemisoides	Feathery Cassia
Ceanothus rigidus	Wild Lilac
Ceratostigma plumbagonoides	
Chamaecyparis pisifera	False Cypress
Chrysanthemum nipponicum	
Cleyera japonica	Cleyera
Citrus spp.	Citrus species*
Coreopsis sp.	Coreopsis (Calliopsis): Early Sunrise, Moonbeam
Cornus florida	Flowering Dogwood
C. stonolifera	American Dogwood
Cortaderia seloana	Pampas Grass
Cotoneaster apiculatus	Cranberry Cotoneaster
C. buxifolius	Cotoneaster
C. Dammeri	Bearberry Cotoneaster

exceed the maximum rate for turfgrass species listed in the MAXIMUM RATE/CALENDAR YEAR of RegalKade 50 by Turfgrass Species Table.

<sup>2</sup>Winter weeds are best controlled by application made prior to germination: late summer; fall or winter timings.

<sup>3</sup>Suppression only. Sequential applications may be made as long as the total amount of product applied does not exceed the maximum annual application rates recommended for the turf species. All applications must be made prior to germination of weed seeds. This product will not control established weeds.

**OVERSEEDING/RESEEDING**

RegalKade 50 will inhibit the germination of turf species if overseeding/ reseeded too soon after application. The interval between application and overseeding/reseeding is dependent on the application rate, soil temperature, and soil moisture. A shorter interval is possible if growing conditions are warm and wet, while a longer interval is needed if conditions are cool and dry.

Use the Overseeding/Reseeding Intervals Table below as a guide when choosing the appropriate overseeding/reseeding interval for each situation.

**OVERSEEDING/RESEEDING INTERVALS TABLE**

RegalKade 50 (lb product/A)	Overseeding/Reseeding Interval (months) Based Upon Seeding Method and Predominant Soil Conditions for the season			
	Broadcast Seeding		Drill Seeding	
	Wet and Warm	Cool and Dry	Wet and Warm	Cool and Dry
100	4	6	3	4
200	8	12	6	8

**LANDSCAPE ORNAMENTALS**

(including Perennial and Wildflower Plantings) RegalKade 50 may be applied for residual pre-emergence weed control in ornamentals.

**USE RATES:** Apply RegalKade 50 at 132 to 300 lb/A in fall and/or spring. Use the higher rate of application for longer control periods. Sequential or single applications are allowed so long as the total amount of product applied does not exceed the maximum annual application rate of 300 lb/A.

**APPLICATION TIMING AND INFORMATION:** Apply RegalKade 50 prior to weed seed germination to the soil surface surrounding established tolerant ornamentals listed below. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply this product after cuttings form roots and become established.

RegalKade 50 is a pre-emergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 in. of rainfall or irrigation, or with shallow (1 to 2 in.) mechanical incorporation, before weed seeds germinate. Acceptable control will be obtained when this product is activated within 14 days after application. Best weed control is obtained when this product is applied to soil free of clods, weeds, and debris such as leaves and mulch. Prior to application of this product, control existing vegetation by hand weeding, cultivation or the use of an appropriate post-emergence herbicide.

to get uniform distribution on the soil surface. RegalKade 50 may be applied over the top of these species. When plants are under stress (such as heat, drought, or frost damage) some cultivars of listed plants may be sensitive to RegalKade 50.

Scientific Name	Common Name
<i>C. microphyllus</i>	Rockspray Cotoneaster
<i>Cretaeus</i> spp.	Hawthorne
<i>Crococsmia</i> sp.	Lucifer
<i>Cupressus sempervirens</i>	Italian Cypress
<i>Delosperma alba</i>	White Trailing Ice Plant
<i>Delosperma</i> sp.	Cooperi Pink
<i>Delphinium</i> sp.	Larkspur: Blue Elf
<i>Dianthus deltooides</i>	Dianthus; Maiden Pinks "Zing"
<i>D. gratianopolitanus</i>	Cheddar Pink
<i>Dodonea viscosa</i>	Hop Bush
<i>Echinacea pupurea</i>	Coneflower, Purple; Mangus
<i>Elaeagnus pungens</i>	Silverberry
<i>Euonymus Fortunei</i>	Wintercreeper
<i>E. japonica</i>	Japanese Spindle Tree (Evergreen Euonymus)
<i>E. kiautschovica</i>	Spreading Euonymus
<i>Fatsia japonica</i>	Japanese Aralia
<i>Forsythia intermedia</i>	Border Forsythia
<i>Forsythia suspensa</i>	Weeping Forsythia
Scientific Name	Common Name
<i>F. viridissima</i>	Greenstem Forsythia
<i>Gaillardia</i> sp.	Gaillardia, Blanket Flower: "Goblin"
<i>Gardenia jasminoides</i>	Gardenia, Cape-Jasmine, Radicans
<i>Gaura</i> sp.	
<i>Gentianna dahurica</i>	Gentian
<i>Geranium Cinereum</i>	Cranesbill
<i>Gladiolus</i> spp.	Gladiolus species**
<i>Gypsophila repens</i>	Baby's Breath
<i>Hedera helix</i>	English Ivy
<i>Hellanthemum</i> sp.	Sunrose
<i>Hemerocallis</i> sp.	Daylily: Aztec Gold, Stella De Oro, Tender Love
<i>Hibiscus</i>	Rose of Sharon**
<i>Hibiscus</i> sp.	Mallow: Disco Belle White
<i>Hibiscus Rosa-sinensis</i>	Chinese Hibiscus
<i>Hosta plantaginea</i>	Hosta, Plantain Lily (Fragrant)
<i>H. Sieboldiana</i>	Hosta, "Searsucker"
<i>Houttuynia cordata</i> var. <i>variegata</i>	
<i>Hydrangea macrophylla</i>	Bigleaf Hydrangea
<i>Ilex Beryi</i>	Holly
<i>Ilex cornuta</i>	Chinese Holly
<i>I. crenata</i>	Japanese Holly: Helleri
<i>I. opaca</i>	American Holly
<i>I. vomitoria</i>	Yaupon Holly: Schillings
<i>Inula ensifolia</i>	
<i>Iris ensata</i>	Sword-Leaved Iris; Jodlesong
<i>Iris siberica</i>	Siberian Iris; Cabernet
<i>Iris</i> spp.	Iris species**
<i>Jasminium nudiflorum</i>	Winter Jasmine
<i>Juniperus chinensis</i>	Chinese Juniper; Nick's Compact, Parsoni
<i>J. conferta</i>	Shore Juniper; Blue Pacific
<i>J. davurica</i>	Parsoni
<i>J. horizontalis</i>	Creeping Juniper
<i>Juglans</i> sp.	Walnut*
<i>Justicia brandegeana</i>	Shrimp Plant

Scientific Name	Common Name
<i>Lagerstromia indica</i>	Crape Myrtle
<i>Lagerstromia indica</i> X <i>Faurei</i>	Crape Myrtle; Tuscarora
<i>Lantana montevidensis</i>	Weeping Lantana
<i>Lavender</i> sp.	Lavender; Munstead
<i>Leontopodium alpinum</i>	Edelweiss
<i>Ligustrum sinense</i>	Chinese Privet; Variegata
<i>Ligustrum amurense</i>	Amur Privet
<i>L. japonicum</i>	Japanese Privet
<i>L. lucidum</i>	Glossy Privet (Waxleaf)
<i>Lillium</i> sp.	Lily; Jazz
<i>Liriope Muscari</i>	Big Blue Lily; Turf; Silver Mound; Evergreen Giant
<i>Liriope Muscari</i> var. <i>variegata</i>	Liriope, Variegated
<i>Liriope spicata</i>	Liriope, Creeping
<i>Lobelia Cardinalis</i>	Cardinal Flower; Indian Pink
<i>Lonicera japonica</i>	Japanese Honeysuckle
Scientific Name	Common Name
<i>L. tatarica</i>	Tatarian Honeysuckle
<i>Loropetalum chinense</i>	Burgundy
<i>Lythrum</i> sp.	Loosestrife; Modern Pink
<i>Magnolia</i> spp.	Magnolia species**
<i>Malephora luteola</i>	Ice Plant
<i>Malus</i> sp.	Crabapple*
<i>Miscanthus sinensis</i>	Yaku jima, Silberfeder
<i>Nandina domestica</i>	Heavenly Bamboo
<i>Narcissus</i> spp.	Narcissus species**
<i>Nerium oleander</i>	Oleander
<i>Olea europea</i>	Olive*
<i>Ophiopogon japonicus</i>	Mondo Grass
<i>Osmanthus heterophyllus</i>	Osmanthus (False Holly): Gulf Tide
<i>Osteospermum fruticosum</i>	Trailing African Daisy
<i>Oxydendrum arboreum</i>	Sourwood
<i>Paeonia suffruticosa</i>	Tree Peony
<i>Pennisetum setaceum</i>	Fountain Grass (Dwarf)
<i>Perovskia atriplicifolia</i>	
<i>Persea americana</i>	Avocado*
<i>Photina Fraseri</i>	Fraser's Photinia (Redtip)
<i>Phytotegia Virgiana</i>	Dragonhead, False; Vivid
<i>Picea</i> spp.	Spruce species** (Colorado Blue, Norway, etc.)
<i>Pieris japonica</i>	Lily-of-the-valley Bush
<i>Pinus brutia</i>	Calabrian Pine
<i>P. canariensis</i>	Canary Island Pine
<i>P. ellioti</i>	Slash Pine
<i>P. halepensis</i>	Aleppo Pine
<i>P. nigris</i>	Austrian Pine
<i>P. palustris</i>	Longleaf Pine
<i>P. radiata</i>	Monterey Pine
<i>P. Thunbergiana J</i>	apanese Black Pine
<i>P. strobus</i>	Eastern White Pine
<i>P. sylvestris</i>	Scotch Pine
<i>P. taeda</i>	Loblolly Pine
<i>P. virginiana</i>	Virginia Pine
<i>Pistachio</i> sp.	Pistachio*

Scientific Name	Common Name
<i>Pittosporum rhombifolium</i>	Queensland Pittosporum
<i>P. tobira</i>	Japanese Pittosporum
<i>Podocarpus macrophyllus</i>	Japanese Yew
<i>Prunus Laurocerasus</i>	English Laurel
<i>Prunus</i> sp.	Almond, Apricot, Nectarine, Peach, Plum and Prune*
<i>Pseudotsuga Menziesii</i>	Douglas Fir**
<i>Pyracantha coccinea</i>	Firethorn (Scarlet)
<i>P. Fortuneana</i>	Firethorn
<i>P. Koidzumii</i>	Forethorn
<i>Pyrus</i> sp.	Bradford Pear sp.
<i>Quercus Shumardii</i>	Oak, Shumard's Red
<i>Quercus</i> spp.	Oak species
<i>Raphiolepis indica</i>	Indian Hawthorne
<i>Ralphiolepis umbellata</i>	Yedda Hawthorn
<i>Rhododendron</i> (including <i>azalea</i> )	'Coral Bells', 'Delaware Valley White', 'Flame Creeper', 'Formosa', 'Girard Crimson', 'George L. Tabor', 'Hi-no-crimson', 'PJM', 'Roseum Elegans', 'Wakeiebsiu', 'White Gumbo'
<i>Rosa Banksiae</i>	Lady Bank's Rose
<i>Rosmarinus officinalis</i>	Rosemary*
<i>Rudbeckia</i> sp.	Black-Eyed Susan; Goldstrum
<i>Santolina virens</i>	Santolina
<i>Saxifraga</i> sp.	Saxifrage; Purple Dome
<i>Scabiosa</i> sp.	Pincushion Flower
<i>Sedum album</i>	Stonecrop
<i>Sedum cauticola</i>	Stonecrop; Lidakense
<i>Sedum dasyphyllum</i>	Stonecrop
<i>Sedum spurium</i>	Stonecrop; Dragon's Blood
<i>Spiraea Bumalda</i>	Spiraea: Anthony Waterer
<i>Syzygium paniculatum</i>	Australian Brushberry
<i>Taxus cuspidata</i>	Japanese Yew
<i>T. media</i>	Yew
<i>Teucrium</i> sp.	Germander
<i>Thalictrum dipterocarpum</i>	Meadow Rue
<i>Thuja occidentalis</i>	American Arborvitae
<i>Trachelospermum asiaticum</i>	Star Jasmine
<i>Tsuga canadensis</i>	Canada Hemlock
<i>Tulipa</i> spp.	Tulip species
<i>Veronica</i> sp.	Veronica, Speedwell; Sunny Border
<i>Viburnum japonicum</i>	Japanese Viburnum
<i>V. odoratissimum</i>	Sweet Viburnum
<i>V. plicatum</i>	Japanese Snowball
<i>V. rigidum</i>	Canary Island Viburnum
<i>V. suspensum</i>	Arrowwood Viburnum
<i>V. tinus</i>	Laurustinus
<i>V. trilobium</i>	Cranberry Bush
<i>V. wrightii</i>	Leatherleaf Viburnum
<i>Vinca major</i>	Greater Periwinkle, Vinca
<i>Vinca minor</i>	Common Periwinkle, Vinca
<i>Vitis</i> sp. Grape*	
<i>Weigela florida</i>	Old Fashioned Weigela
<i>Yucca aloifolia</i>	Spanish Bayonet, Yucca
<i>Y. filamentosa</i>	Adam's Needle, Yucca

\*Ornamental species only. Do not use on food producing trees or vines.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original container away from other fertilizer, feed, or food stuffs and separated from other pesticides.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**STORAGE:** Store in original container away from other fertilizer, feed, or food stuffs and separated from other pesticides.

**DISPOSAL:** If empty: **DO NOT** reuse this container. Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

#### CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once. The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of this product, which are beyond the control of REGAL CHEMICAL COMPANY, Inc. or Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer and User; and Buyer and User agree to hold REGAL and Seller harmless for any claims relating to such factors.

REGAL warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or REGAL, and Buyer and User assume the risk of any such use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, REGAL MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither REGAL or Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF REGAL AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF REGAL OR SELLER, THE REPLACEMENT OF THE PRODUCT.

REGAL and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of REGAL.

