

**RESTRICTED USE PESTICIDE**

Toxic to fish and aquatic organisms

For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicator's certification.



**Arctic<sup>®</sup> 3.2EC**

**Insecticide**

For Agricultural or Commercial Use Only

**For Control of Soybean Aphid in Soybeans**  
**In All States Where Crop and Pest Exist and Where 2(ee) Recommendations Are Recognized.**

**2(ee) Recommendation**

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

**Directions For Use:**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

CROP	PEST	RATE OF APPLICATION
Soybeans	Soybean Aphid	4 to 8 ounces (0.1 to 0.2 lb. active) per acre

Use Arctic 3.2EC at the rate of 4 to 8 ounces (0.1 to 0.2 pound a.i.) per acre. For better coverage use with a spreader type adjuvant such as Silkin<sup>™</sup> or Class Act<sup>®</sup> Next Generation. For better coverage and increased residual control use with a spreader type adjuvant such Preference<sup>®</sup>. Use of higher volumes of carrier will also result in more complete plant coverage and better control. Use the higher recommended dosage for increased pest pressure and for increased residual control.

Use Restrictions: Do not apply more than 0.4 pound active ingredient per acre per season. Do not apply within 60 days of harvest. Do not graze or feed soybean forage or hay.

**IMPORTANT:**

**All applicable directions, restrictions and precautions on the EPA registered label are to be followed.**

**Personal protective equipment and reentry intervals on the basic label must be followed.**

Distributed By:

Agrilience, LLC, P.O. Box 64089, St. Paul, MN 55164-0089

Arctic<sup>®</sup> is a registered trademark of FMC Corporation