

FIFRA Section 2(ee) Recommendation



GROUP	14	HERBICIDE
-------	----	-----------

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF
AL, AR, DE, GA, IA, IL, IN, KS, KY, LA, MD, MI, MO, NC, NE,
NJ, OH, OK, SC, TN, VA and WI



EPA Reg. No. 59639-117

**VALOR® XLT SOYBEAN HERBICIDE PLUS BOUNDARY®, ALACHLOR,
S-METOLACHLOR, DIMETHENAMID-P OR FLUFENACET HERBICIDES FOR EARLY
PRE-PLANT APPLICATIONS AT A MINIMUM OF 14 DAYS PRIOR TO PLANTING FOR
RESIDUAL CONTROL OF CERTAIN WEEDS IN NO-TILLAGE AND REDUCED-TILLAGE
SOYBEANS**

OR

**VALOR XLT SOYBEAN HERBICIDE APPLIED AT A MINIMUM OF 14 DAYS PRIOR TO
PLANTING FOLLOWED BY BOUNDARY, ALACHLOR, S-METOLACHLOR,
DIMETHENAMID-P OR FLUFENACET HERBICIDES APPLIED AT PLANTING FOR
RESIDUAL CONTROL OF CERTAIN WEEDS IN NO-TILLAGE AND REDUCED-TILLAGE
SOYBEANS**

THIS FIFRA SECTION 2(EE) EXPIRES DECEMBER 31, 2021

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This 2(ee) recommendation contains directions for use which do not appear on the package label and must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions and precautions on the EPA registered label when using *Valor* XLT Soybean Herbicide. Always read and follow all label directions when using any pesticide alone or in tank mix combinations. The most restrictive labeling applies when using a tank mix.

- In the states of Delaware, Maryland, New Jersey and Virginia do not exceed 3.0 oz/A of *Valor* XLT Soybean Herbicide.
- Do not graze animals on green forage or stubble. Do not utilize hay or straw for animal feed or bedding.

WEEDS CONTROLLED	ORGANIC MATTER	SOIL TYPE	RATES
Lambsquarters, Common Nightshades Black Eastern Black Pigweeds Redroot Palmer Amaranth Smooth Spiny Amaranth Tumble Sida, Prickly Velvetleaf ¹	Up to 5%	All Soil Types	<i>Valor</i> XLT Soybean Herbicide at 3 to 5 oz/A plus Boundary 7.8 EC at 0.5 to 2 pt/A or Boundary 6.5 EC at 0.6 to 2.4 pt/A or alachlor at 1 to 2 lb ai/A or s-metolachlor at 0.5 to 2 lb ai/A or dimethenamid-p at 0.25 to 0.6 lb ai/A or flufenacet at 0.15 to 0.45 lb ai/A
APPLICATION METHOD: These early preplant applications must be surface applied. Incorporation of these tank mixes will result in reduced control of the weeds listed above. NOTE: Refer to <i>Valor</i> XLT Soybean Herbicide, Boundary 7.8 EC, Boundary 6.5 EC, alachlor, s-metolachlor, dimethenamid-p or flufenacet labels for other weeds controlled.			

¹A postemergence herbicide, such as *Resource*[®] or glyphosate plus *Resource* (Roundup Ready[®] soybeans only) may be needed to adequately control velvetleaf.

Please contact Valent U.S.A. Corporation at 800-6-VALENT (682-5368) to determine if this product is registered in your state.

©2016 Valent U.S.A. Corporation

Resource and *Valor* are registered trademarks of Valent U.S.A. Corporation.
Boundary is a registered trademark of Syngenta, EPA Reg. No. 100-1162 264-771
Roundup Ready is a registered trademark of Monsanto Company.

Manufactured for:
Valent U.S.A. Corporation
P.O. Box 8025
Walnut Creek, CA 94596-8025

Made in U.S.A.