

RESTRICTED USE PESTICIDE

Toxic to fish and aquatic organisms

For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicator's certification.



Arctic[®] 3.2EC
Insecticide

For Agricultural or Commercial Use Only

For Control of Southern Corn Rootworm in Corn

2(ee) Recommendation

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

Directions For Use:

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

CROP	PEST	RATE OF APPLICATION
Corn (field), Field Corn Grown for Seed, Popcorn	Southern Corn Rootworm	4 to 8 ounces (0.1 to 0.2 lb. active) per acre as a T-band application at planting.

Preemergent Use: To control Southern Corn Rootworm, use Arctic 3.2 EC at a rate of 4 to 8 ounces per acre at planting time. For best results, apply as an in-furrow or T-band treatment. This can also be done as a split application of 2 oz. per acre banded with a pop up fertilizer plus 2 oz. per acre banded with a preemergent herbicide at planting. Be sure that the herbicide label does not prohibit application with insecticides and to read and follow all precautionary statements on the herbicide label. When using Arctic 3.2EC and liquid pop-up fertilizers, a jar compatibility test should be performed with appropriate ratio of both products to ensure mixture will stay in solution. Constant agitation should be maintained during mixing and application.

IMPORTANT: All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

Personal protective equipment and reentry intervals on the basic label must be followed.

Distributed By:

Agrilience, LLC, P.O. Box 64089, St. Paul, MN 55164-0089

Arctic[®] is a registered trademark of FMC Corporation

0/A20/2