



Bayer CropScience

**Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)**

SCALA BRAND SC FUNGICIDE

EPA Reg. No. 264-788

For Use On: Bulb Vegetables, Lemon, Small Fruit vine climbing, Stone Fruit, Pome Fruit, Low growing berry crop subgroup, Tomato

This supplemental label expires on 11/20/2018 and must not be used or distributed after this date.

Supplemental Label

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Read this label and the product package label before using this product. This Supplemental Label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the product label for SCALA BRAND SC FUNGICIDE attached to the container.

BULB VEGETABLES CROP GROUP 3-07

CROP	DISEASE	USE RATE		USE INFORMATION
		FL OZ PER ACRE	LBS A.I. PER ACRE	
Onion Bulb Subgroup Daylily, bulb; fritillaria, bulb; garlic, bulb; garlic, great-headed, bulb; garlic, serpent, bulb; lily, bulb; onion, bulb; onion, Chinese, bulb;	Botrytis leaf blight, Neck rot (<i>B. squamosa</i> <i>B. allii</i>) Purple blotch (<i>Alternaria porri</i>)	SCALA BRAND SC FUNGICIDE Alone		When using tank mix rates, add fungicide(s) registered for use against the targeted disease(s). Use lower rate only in a tank mix with a broad spectrum fungicide for bulb vegetables. For optimum results, begin applications as soon as crop and/or environmental conditions become favorable for disease development. Make applications on a 7 to
		18	0.70	
		SCALA BRAND SC FUNGICIDE In tank mixes		

<p>onion, pearl; onion, potato, bulb; shallot, bulb; cultivars, varieties, and/or hybrids of these.</p> <p>Onion Green Subgroup Chive, fresh leaves; chive, Chinese, fresh leaves; elegans hosta; fritillaria, leaves; kurrat; lady's leek; leek; leek, wild; Onion, Beltsville bunching; onion, fresh; onion, green; onion, macrostem; onion, tree, tops; onion, Welsh, tops; shallot, fresh leaves; cultivars, varieties, and/or hybrids of these.</p>		9 - 18	0.35 - 0.70	<p>14 day interval depending upon disease conditions.</p> <p>RESTRICTIONS: Do not apply more than 54 fl oz/A (2.1 lbs a.i./A) of SCALA BRAND SC FUNGICIDE per crop. Do not apply within 7 days of harvest.</p>
--	--	--------	-------------	---

LEMONS

CROP	DISEASE	USE RATE		USE INFORMATION
		FL OZ PER ACRE	LBS A.I. PER ACRE	
<p>Australian desert lime; Australian finger-lime; Australian round lime; brown river finger lime; kumquat; lemon; lime; mount white lime; New Guinea wild lime; Russell River lime; sweet lime; Tahiti lime; cultivars, varieties, and/or hybrids of these</p>	<p>Alternaria (<i>Alternaria</i> spp. - Suppression only)</p> <p>Penicillium rot (<i>Penicillium</i> spp. - Suppression only)</p>	18	0.70	<p>Always consult your agricultural advisor, University contact or Extension Service for recommended pest management practices for your area.</p> <p>The use of adjuvants is prohibited.</p> <p>RESTRICTIONS: Make one application at the rate of 18 fl oz/A (0.70 lbs a.i./A). Do not apply more than 18 fl oz/A (0.70 lbs a.i./A) of SCALA BRAND SC FUNGICIDE per year. Do not apply within 7 days of harvest.</p>

SMALL FRUIT VINE CLIMBING CROP SUBGROUP 13-07 F EXCEPT FUZZY KIWIFRUIT

CROP	DISEASE	USE RATE		USE INFORMATION
		FL OZ PER ACRE	LBS A.I. PER ACRE	
Amur river grape; gooseberry; grape; kiwifruit, hardy; maypop; schisandra berry; cultivars, varieties, and/or hybrids of these	Gray mold (<i>Botrytis cinerea</i>)	SCALA BRAND SC FUNGICIDE Alone		When using tank mix rates, add fungicide(s) registered for use against the targeted disease(s). Use lower rate only in a tank mix with another fungicide active against <i>Botrytis</i> . Apply product at the critical timings for <i>Botrytis</i> control. Typically, first applications are made at early bloom, and/or berry touch to bunch closure, veraison, and pre-harvest. Use sufficient water to ensure penetration of the canopy and coverage of the flowers or bunches. RESTRICTIONS: Do not apply more than 36 fl oz/A (1.4 lbs a.i./A) of SCALA BRAND SC FUNGICIDE per crop. Do not apply within 7 days of harvest.
		18	0.70	
		SCALA BRAND SC FUNGICIDE In tank mixes		
		9 - 18	0.35 - 0.70	

STONE FRUITS GROUP 12-12 (Except Cherries)

CROP	DISEASE	USE RATE		USE INFORMATION
		FL OZ PER ACRE	LBS A.I. PER ACRE	
Apricot; apricot, Japanese; capulin; Jujube, Chinese; nectarine; peach; plum; plum, American; plum, beach; plum, Canada; plum, cherry; plum, Chickasaw; plum, Damson; plum, Japanese; plum, Klamath; plum, prune; plumcot; sloe; cultivars, varieties, and/or hybrids of these	Brown rot blossom blight (<i>Monilinia</i> ssp.)	SCALA BRAND SC FUNGICIDE Alone		For optimum results, begin applications as soon as crop and/or environmental conditions become favorable for disease development. Apply when bud tissue is susceptible to infection (i.e., pink, white or red bud). If conditions favorable for disease development persist or recur, apply at full bloom or at petal fall. Make pre-harvest applications on 7 day intervals or longer. When using tank mix rates, add fungicide(s) registered for use against the targeted disease(s). RESTRICTIONS: Do not apply more than three applications of SCALA BRAND SC FUNGICIDE alone or more than 54 fl oz/A (2.1 lbs a.i./A) of SCALA BRAND SC FUNGICIDE per year. Do not apply within 2 days of fruit harvest or make more than 2 applications of a Group 9 fungicide within 30 days of harvest. Do not use on cherries.
		9 - 18	0.35 - 0.70	
	Jacket rot, Green fruit rot (<i>Botrytis cinerea</i>)	SCALA BRAND SC FUNGICIDE Alone		
		18	0.70	
	Shot hole (<i>Wilsonomyces carpophilus</i>)	SCALA BRAND SC FUNGICIDE In Tank Mixes		
		9 - 18	0.35 - 0.70	
	Brown fruit rot (<i>Monilinia fructicola</i>)			

POME FRUITS GROUP 11-10

CROP	DISEASE	USE RATE		USE INFORMATION
		FL OZ PER ACRE	LBS A.I. PER ACRE	
Apple; azarole; crabapple; loquat; mayhaw; medlar; pear; pear, Asian; quince; quince, Chinese; quince, Japanese; tejocote; cultivars, varieties, and/or hybrids of these	Scab (<i>Venturia</i> spp.)	SCALA BRAND SC FUNGICIDE Alone		Under conditions favorable for disease development, shorten the spray intervals and/or use the high rate. When using tank mix rates, add fungicide(s) registered for use against the targeted disease(s). For optimum results, begin applications at green tip or as soon as crop and/or environmental conditions become favorable for disease development. Make pre-harvest applications on 7 day intervals or longer. RESTRICTIONS: Do not apply more than 40 fl oz/A (1.6 lbs a.i./A) of SCALA BRAND SC FUNGICIDE per year. Do not apply within 72 days of harvest.
		7 - 10	0.27 - 0.39	
		SCALA BRAND SC FUNGICIDE In tank mixes		
		5 - 10	0.19 - 0.39	

LOW GROWING BERRY CROP SUBGROUP 13-07 G

CROP	DISEASE	USE RATE		USE INFORMATION
		FL OZ PER ACRE	LBS A.I. PER ACRE	
Bearberry; bilberry; blueberry, lowbush; cloudberry; cranberry; lingonberry; muntries; partridgeberry; strawberry; cultivars, varieties, and/or hybrids of these	Gray mold (<i>Botrytis cinerea</i>)	SCALA BRAND SC FUNGICIDE Alone		When using tank mix rates, add fungicide(s) registered for use against the targeted disease(s). For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. Make applications from pre-bloom to harvest on a 7 to 14 day interval depending upon disease conditions. Always follow resistance management directions listed on this label and the recommendations for your area. RESTRICTIONS: Do not apply more than 54 fl oz/A (2.1 lbs a.i./A) of SCALA BRAND SC FUNGICIDE per crop. Do not apply within 1 day of harvest.
		18	0.70	
		SCALA BRAND SC FUNGICIDE In tank mixes		
		9 - 18	0.35 - 0.70	

TOMATOES SUBGROUP 8-10A

CROP	DISEASE	USE RATE		USE INFORMATION
		FL OZ PER ACRE	LBS A.I. PER ACRE	
Bush tomato; cocona; currant tomato; garden huckleberry; goji berry; groundcherry; naranjilla; sunberry; tomatillo; tomato; tree tomato; cultivars, varieties, and/or hybrids of these	<p>Early blight (<i>Alternaria solani</i>)</p> <p>Gray mold (<i>Botrytis cinerea</i>)</p>	7	0.27	<p>Under conditions favorable for disease development, shorten the spray intervals and/or increase the rate of the tank-mix partner.</p> <p>Use only in a tank mix with labeled dose rate of another effective early blight fungicide. For optimum results, begin applications as soon as crop and/or environmental conditions become favorable for disease development. Make applications on a 7 to 14 day interval depending upon disease conditions.</p> <p>RESTRICTIONS:</p> <p>Do not apply more than 35 fl oz/A (1.4 lbs a.i./A) of SCALA BRAND SC FUNGICIDE per crop, with a maximum of 2 crop cycles per year. Do not apply within 1 day of harvest.</p> <p>Greenhouse Use: Apply SCALA BRAND SC FUNGICIDE only in well-ventilated plastic tunnel houses or glass houses. Ventilate for at least 2 hours after application.</p>

**For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577
For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)**

*As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*