

FIFRA Section 24(c) Special Local Need (SLN) Label

Corteva Agriscience

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Starane[®] Ultra

EPA Reg. No. 62719-577

EPA 24(c) Special Local Need Registration SLN DE-150001
For Distribution and Use Only Within the State of Delaware

This label expires and must not be distributed or used in accordance with this SLN registration after July 1, 2030.

90-Day Plantback Restriction to Soybeans After a Spring Application to Winter Wheat to Control Broadleaf Weeds Including Chickweed

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Starane[®] Ultra herbicide before applying. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.
- Use of Starane Ultra according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Starane Ultra.

Directions for Use

Apply as a broadcast postemergence treatment to actively growing wheat from the 2 leaf crop growth stage up to and including flag leaf emergence (Zadoks scale 39) for control of broadleaf weeds including chickweed. Apply when weeds are actively growing, but before weeds are 8 inches tall or vining. Only weeds emerged at the time of treatment will be controlled. Extreme growing conditions such as drought or near freezing temperatures prior to, at, and following time of application may reduce weed control and increase the risk of crop injury at all stages of growth. **Do not use if cereal crop is underseeded with a legume.**

Spot Application: Spot applications may be made; however, to prevent over application, apply spot treatments at rates and spray volumes equivalent to broadcast application. See instructions for Spot Application in Application Directions section on the label affixed to the product.

Broadcast Application Rates:

Weed Size or Species	Application Rate (pint/acre)
susceptible broadleaf weed seedlings less than 4 inches tall	0.3
susceptible broadleaf weed seedlings less than 8 inches tall or vining	0.4
volunteer potatoes	0.7

Restrictions:

- **Preharvest Interval:** Do not apply closer than 14 days before cutting of hay or 40 days before harvesting of grain and straw.
- Do not allow livestock to graze treated areas or harvest treated forage within 7 days of application.
- Do not apply more than 0.7 pint of Starane Ultra per acre per growing season.

- **Plantback Restriction:** 90 days must have elapsed from time of the last application of Starane Ultra until the planting of soybeans. For all other plantback restrictions, see the label affixed to the container.

^{TM®} Trademarks of Corteva Agriscience and its affiliated companies

R357-063

Accepted: 07/25/2025

Replaces: R357-039