



Bayer Environmental Science, a division of Bayer
CropScience
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-800-331-2867

Tartan® Stressgard®

EPA Reg. No. 432-1446

For control of Take-All Root Rot,
Bermudagrass decline, and Warm Season
Turfgrass Decline (*Gaeumannomyces*
graminis var. *graminis*)

Expires: 6/14/2020

PRODUCT BULLETIN

**FOR DISTRIBUTION AND USE ONLY IN THE STATES OF FLORIDA, TEXAS,
HAWAII, GEORGIA, SOUTH CAROLINA, NORTH CAROLINA, TENNESSEE,
ARKANSAS, OKLAHOMA, KANSAS, MISSOURI, MISSISSIPPI, ALABAMA,
LOUISIANA, NEW MEXICO, AND ARIZONA**

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. It is a violation of federal law to use this product in a manner inconsistent with its label.

This product bulletin must be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

For use on turfgrass on golf courses, sod farms and sports fields only.

Disease	Tartan® Stressgard®	fl oz per 1,000 sq ft	Application Interval/Timing
Take-All Root Rot,	Apply 1 to 2 preventive applications in the spring and fall, prior to conditions favorable for disease development.	2 fl oz	28 days
Bermudagrass Decline,		2 fl oz	28 days
Warm Season Turfgrass Decline (<i>Gaeumannomyces graminis</i> var. <i>graminis</i>)	Apply fungicide in adequate water volume and water-in applications to root zone.	2 fl oz	28 days

For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-800-331-2867

As with any plant-protection product, always read and follow label directions.