



FIRESTORM[®]

**Defoliant and desiccant herbicide
for the control of weeds and grasses
and as a harvest aid**

PRODUCT INFORMATION BULLETIN

Supplemental Recommendation

**For the Use of FIRESTORM – EPA Reg. No. 82557-1-400
on SOYBEANS**

**FIFRA 2(ee) recommendation for use and distribution in
the state of Louisiana only.**

**Under FIFRA 2(ee) FIRESTORM is recommended
for use as a harvest aid for soybeans**

Use Pattern	FIRESTORM Rate/A	Minimum Total Spray per Acre	Grazing or Preharvest Interval (Days)	Additional Precautions, Restrictions and Directions
Harvest Aid	5.4-10.7 fl. oz.	Ground: 10 gals. Air: 5 gals.	--	<ul style="list-style-type: none">• Do not make more than 3 applications per year.• Indeterminant varieties: Applications should be made when at least 65% of the seed pods have reached a mature brown color or when seed moisture is 30% or less.• Determinant varieties: Apply when plants are mature, i.e., beans are fully developed, 1/2 of leaves have dropped, and remaining leaves are yellowing.• Injury will occur on immature soybeans.• Mature cocklebur, especially drought-stressed plants, are tolerant to FIRESTORM and desiccation will not be complete. Always use the higher rate when treating cocklebur.• Do not apply within 15 days of harvest.• Do not graze or harvest for forage or hay.

This recommendation is made as permitted under FIFRA section 2(ee) and has not been submitted to or approved by the EPA. **Observe the warranty statements and conditions specified on the label for FIRESTORM. It is important to read and observe all restrictions, precautions and other information appearing on the label.**

FIRESTORM[®] is a registered trademark of Chemtura Corporation

08-17-2012