Product Bulletin

Matrix® SG

EPA Reg. No. 352-768

Special 2(ee) Recommendation[†]

For Distribution and Use Only in the States of Arizona, California, Idaho, Utah and Washington

MATRIX® SG PLUS "GOAL 2XL" OR "GOALTENDER" OR "CHATEAU SW" FOR IMPROVED WEED CONTROL IN CITRUS, GRAPES, POME FRUITS, STONE FRUITS AND TREE NUTS

ATTENTION

[†]This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Matrix SG herbicide before applying. Carefully follow all
 precautionary statements and applicable use directions.
- Except as specified in this product bulletin, use of these products is subject to all use precautions and limitations imposed by the label affixed to the container.

DIRECTIONS FOR USE

To the extent consistent with applicable law, DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with uses not specified by the label or 2(ee) instructions.

MATRIX® SG Herbicide applied at the rate of 4.0 oz product (0.0625 lb ai) per acre can be tank mixed with Goal ® 2XL (Al: oxyfluorfen, Reg No. 62719-424) at 3.0 to 6.0 pints (0.75 - 1.5 lb ai) per acre, or GoalTender® herbicide (Al: oxyfluorfen, Reg. No. 62719-447) at 1.5 to 3.0 pints (0.75 - 1.5 lb ai) per acre, or "Chateau SW" herbicide (Al: flumioxazin, EPA Reg No. 59639-99) at 6.0 to 12.0 ounces (0.2 - 0.4 lb ai) per acre, to help improve overall weed control in citrus, grapes, pome fruits, stone fruits and tree nuts. For best results incorporate mechanically via light tillage or with 1/2" irrigation following application.

It is important to use a postemergence/contact herbicide including glyphosate or other registered postemergence herbicides to control any emerged weeds in order to obtain the best residual weed control from MATRIX® SG Herbicide.

It is the pesticide user's responsibility to ensure that all product are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Read and follow all mixing instructions on the MATRIX® SG label.

- **Do not** apply more than 4.0 oz (0.0625 lb ai) MATRIX® SG Herbicide per year on a broadcast acre basis.
- When applied as a banded treatment (50% band or less), MATRIX® SG Herbicide may be applied twice per year.
- Allow a minimum of 30 days between applications of MATRIX® SG Herbicide.
- Weed control is limited to weeds on the MATRIX® SG Herbicide label.

®™ Trademarks of Dow AgroSciences, DuPont or Pioneer and their affiliated companies or respective owners.

"Chateau" is a registered trademark of Valent USA LLC

E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Rd, Wilmington, DE 19801 U.S.A.

R628-006

Issued: 04/21/2020 Replaces: R628-001