

SUPPLEMENTAL USE DIRECTIONS

In Season Applications on Grapes, Berries, Stone Fruit, Pome Fruit, Citrus



FOR THE PREVENTION AND CONTROL OF POST HARVEST DISEASE

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUCIÓN

See additional precautionary statements elsewhere on this label.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

This product has not been registered by the U.S. Environmental Protection Agency (EPA). SEIPASA SA represents that this product qualifies for exemption from registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Crop	Pest	Rate (Oz/100 gallons)	Dilution (Gal/Acre)	Timing/Application
Stone Fruit Pome Fruit Berries Grapes Citrus Tropical Fruit	Anthracnose (<i>Colletotrichum</i> spp.) Black Mold (<i>Aspergillus niger</i>) Blue mold (<i>Penicillium</i> spp.) Brown Rot (<i>Monilinia</i> spp.) Bulls Eye Rot (<i>Neofabraea</i> spp.) Green mold (<i>Penicillium digitatum</i>) Grey mold (<i>Botrytis cinerea</i>) Mucor Fruit Rot (<i>Mucor</i> spp.) Phomopsis Soft Rot (<i>Phomopsis</i> spp.) Rhizopus Fruit Rot (<i>Rhizopus</i> spp.) Stem End Rot (<i>Gnomonia comari</i>) Sour Rot (<i>Geotrichum</i>)	32-50	50 – 300 by ground	Thorough spray coverage is essential for good pest control. Preharvest applications should be made 7-10 days prior to harvest In season applications can be made for sanitation / prevention.

Manufactured by:

Seipasa, S. A
22240 Tardienta (Huessa)
Spain
www.seipasa.com

Distributed by:

Sym-Agro, Inc
111 South Court Street, Ste 206
Visalia, CA 93291
www.sym-agro.com