

Product Bulletin



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Enable[®] 2F

EPA Reg. No. 62719-416

Special 2(ee) Recommendation[†]

Control of Postbloom Fruit Drop (*Colletotrichum acutatum*) of Citrus Fruits (For Distribution and Use Only in the State of FL)

ATTENTION

[†]This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Enable[®] 2F fungicide before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Use of Enable 2F according to this Section 2(ee) recommendation is subject to all use precautions and limitations imposed by the label affixed to the container for Enable 2F.

Directions for Use

Pest	Enable [®] 2F (fl oz/acre)
Postbloom Fruit Drop (PFD) (<i>Colletotrichum acutatum</i> ¹)	8

¹Regularly monitor disease incidence and weather to initiate fungicide applications.

Refer to product label for General Use Precautions, Mixing Directions, Application Directions, and Restrictions.

Application Timing: Make the first protectant application at the recommended timing for Postbloom Fruit Drop (PFD) of citrus fruits in your area. Decisions are based on previous history of PFD and weather conditions. Treat in accordance with local economic thresholds. Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

Resistance Management: Enable 2F belongs to the sterol demethylation inhibitor (DMI) class of fungicides (FRAC Group 3). Since certain fungi can develop resistance to this class of products, use Enable 2F as part of a resistance management strategy that includes alternation and/or mixing with fungicides of different target site of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

Restrictions:

- Do not make more than 3 applications or apply more than 24 fl oz of Enable 2F (0.38 lb active) per acre per year.
- **Preharvest Interval:** Up to day of harvest.

- **Minimum Treatment Interval:** 21 Days.
- Do not graze livestock in treated areas or feed cover crops grown in treated areas of livestock.
- Do not apply Enable 2F with polymer based spray adjuvants.
- Do not apply this product by chemigation to citrus.

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

Issued: 12/12/17
Initial printing