

SUPPLEMENTAL USE DIRECTIONS

Expanded Use Rates for Vegetable and Herb Crops



TO AID IN THE CONTROL OF DIFFICULT PESTS

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUCIÓN

See additional precautionary statements elsewhere on this label.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

This product has not been registered by the U.S. Environmental Protection Agency (EPA). SEIPASA SA represents that this product qualifies for exemption from registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Crops including but not limited to	Pest	Rate (Oz/100 gallons)	Dilution (Gal/Acre)	Timing/Application
Vegetables: Asparagus, Bean, Beet, Broccoli, Brussels sprouts, Cabbage, Carrot, Cauliflower, Celery, Cucumber, Eggplant, Endive, Garlic, Leek, Lettuce, Melon, Okra, Onion, Pepper, Potato, Pumpkin, Radicchio, Shallots, Spinach, Squash, Swiss chard, Tomato Herbs: Alfalfa, Basil, Chives, Cilantro, Dill, Hops, Mint, Oregano, Parsley, Thyme, Hemp, Cannabis	Aphids (such as cotton aphid, green peach aphid among others). Thrips (such as Thrips spp., Frankliniella spp., Heliothrips spp., Scirtothrips citri among others.) Leafhoppers (such as Empoasca spp., Macrosteles spp., and Graphocephala spp.) White flies (such as Dialeurodes spp., Trialeurodes spp., Bemisia spp.) Armyworms, Caterpillars and loopers (such as cabbage looper, tent caterpillar, fruit worms, alfalfa caterpillar, caterpillar larvae among others)	32-64	40 – 200 by ground	Thorough spray coverage is essential for good pest control. Heavy population management can require sequential treatments on a 5-7-day spray interval. Test Sensitive crops before using the highest rates to ensure acceptable crop safety Do not apply to wilted or otherwise stressed plants, or to newly transplanted material prior to root establishment.

Manufactured by:

Seipasa, S. A
22240 Tardienta (Huesca)
Spain
www.seipasa.com

Distributed by:

Sym-Agro, Inc
111 South Court Street, Ste 206
Visalia, CA 93291
www.sym-agro.com