

CAUTION: KEEP OUT OF REACH OF CHILDREN

HAZARD STATEMENTS: May be harmful if swallowed. Causes mild skin irritation. Causes eye irritation. May be harmful if inhaled.

FIRST AID: In case of eye contact: Rinse eyes with plenty of water for 15 minutes. Remove contact lenses if present and easy to do. Get medical attention if irritation persists. **In case of skin contact:** Rinse skin with plenty of water for 15 minutes. Take off contaminated clothing and shoes. Get medical attention if irritation persists after rinsing. **If swallowed:** Rinse mouth. Do NOT induce vomiting. If vomiting occurs, keep head low so that stomach content doesn't get into the lungs. **If inhaled:** Move person to fresh air. If symptoms persist after rinsing, get medical attention. If breathing stops, administer artificial respiration.

ENVIRONMENTAL PRECAUTIONS: This product may be toxic to fish and other aquatic organisms. For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Not for aquatic use.

STORAGE AND DISPOSAL: Store this product in its original container in a cool, dry, well-ventilated area, out of reach of children and animals. Dispose of contents and empty package in accordance with municipal regulations. Never discard any unused product in any indoor or outdoor drain.

"TRIPLE WASH EMPTY CONTAINERS AND POUR THE RINSE WATER INTO A CONTAINER THAT IS USED TO PREPARE THE MIXTURE."

WARRANTY: Ultraquimia Agrícola, S.A. de C.V., guarantees the net content and composition of the product published on this label but is not responsible for the seal on the container being broken or damaged and does not guarantee or take responsibility for the improper use and handling during transport or storage application.

RESISTANCE MANAGEMENT

"TO PREVENT THE DEVELOPMENT OF RESISTANT POPULATIONS, ALWAYS RESPECT THE DOSE AND APPLICATION INTERVALS ON THIS LABEL: AVOID REPEATED USE OF THIS PRODUCT; ALTERNATE WITH OTHER CHEMICAL GROUPS WITH DIFFERENT MODES OF ACTION AND DIFFERENT MECHANISMS OF DETOXIFICATION; USE OTHER METHODS OF CONTROL."

MADE IN MEXICO

For use in Organic Production



Cinnamon Oil INSECTICIDE & MITICIDE / OIL IN WATER EMULSION

PERCENT COMPOSITION	BY WEIGHT
ACTIVE INGREDIENT	
Cinnamon oil.	4.00%
INERT INGREDIENTS	
Water, Soap, Xanthan gum.	96.00%
	Total: 100.00%

This product has not been registered by the United States Environmental Protection Agency (EPA). Ultraquimia Agrícola, S.A. de C.V., maintains that this product qualifies for exemption from registration under the Federal Insecticide, Fungicide and Rodenticide Act, under section 25(b).

Net Contents:	32 fl. oz. (946 mL)	12 containers of 32 fl. oz.
	1 Gal (3.78 L)	4 containers of 1 Gal
	2.5 Gal (9.46 L)	2 containers of 2.5 Gal

Keep Out of Reach of Children

CAUTION
See back for additional precautionary statements

Lot #:
Date of Manufacture:

Expiration Date:
2 years, if stored in
conditions described
on the label

MANUFACTURED BY:
Ultraquimia Agrícola, S.A. de C.V.
56 Sur, N 7 A (Esq. Andador 11),
Col. CIVAC, 62500, Jiutepec, Morelos, Mexico
Tel. (777) 3211477.



US Customer Service
Toll Free Number:
1(800) 801-9657

DIRECTIONS FOR USE

Shake the product before using. Adjust the pH of the water to between 5.5 and 6. It is recommended that applications be performed when the first ovipositions are observed or when the first infestation is observed, preferably the first nymphal instars which are the most susceptible. Ensure that the product contacts the insects and that the size of the drop and the pressure allow coverage of the areas where the pest is located. Apply higher doses to serious infestations. **Restrictions:** apply the solution the same day it is prepared. Do not apply to plants that are stressed by inadequate irrigation, in extreme temperatures, when the sun is strong, when it is raining or when it is windy and can cause drift. CinnAcar is not phytotoxic to the crops and in the dose recommended here and if it is applied in accordance with the recommendations on this label. **Incompatibility:** it is not compatible with alkaline or oxidizing substances. Do not mix with products whose compatibility is not completely tested.

USE ONLY ON THE CROPS AND FOR THE PESTS RECOMMENDED HERE

CinnAcar is a botanical insecticide and miticide that may be used according to the following recommendations:

Crop	Pest	Rate (fl.oz./100 gallons)	Gallons /Acre	Remarks
Avocado (NL)	Red spider mite (<i>Oligonychus punicae</i>)	13 - 26	160 - 214	Make 3 applications to foliage at 7-day intervals.
Cotton (NL)	Red spider mite (<i>Tetranychus urticae</i>) Whitefly (<i>Bemisia tabaci</i>) Thrips (<i>Frankliniella occidentalis</i>)	26 - 51	21 - 53	Make 3 applications to foliage at 7-day intervals.
Garlic, Onion, Chives, Asparagus (NL)	Thrips (<i>Thrips tabaci</i>)	28 - 57	48	Make 2 applications to foliage at 8-day intervals.
Asparagus (NL)	European Asparagus Aphid (<i>Brachycorynella asparagi</i>)	35 - 45	25 - 100	Make 3 applications to foliage at 7-day intervals.
Cranberry, Strawberry, Current, Blackberry (SL)	Red spider mite (<i>Tetranychus urticae</i>) Thrips (<i>Frankliniella occidentalis</i>)	43 - 64	64	Make 3 applications to foliage at 7-day intervals.
Blueberry (NL)	Mealybug (<i>Dysmicoccus vaccinii</i>)	14 - 30	91 - 101	
Eggplant, Chili, Pepper, Tomato, Potato, Bell pepper, Tobacco, Tomatillo (NL)	Whitefly (<i>Bemisia tabaci</i>)	45 - 78	35 - 46	Make 3 applications to foliage, at 7-day intervals, beginning 20 days after transplanting.
Summer squash, Winter Squash, Chayote, Melon, Cucumber, Watermelon (NL)	Whitefly (<i>Bemisia tabaci</i>)	30 - 47	59 - 69	Make 3 applications to foliage, at 7-day intervals.
Lime, Lemon, Mandarin, Orange, Tangerine, Grapefruit (NL)	Asian citrus psyllid (<i>Diaphorina citri</i>) Broad mite (<i>Polyphagotarsonemus latus</i>) Citrus mealybug (<i>Pianococcus citri</i>)	26 - 77	53	Make 5 applications to foliage at 21-day intervals.
Prickly Pear cactus (NL)	Cochineal (<i>Dactylopius coccus</i>)	23	182	Make 3 applications to foliage at 7-day intervals.
Grape (NL)	Aphids (<i>Aphis illinoisensis</i>)	22 - 78	52 - 63	Start applications when the first infestation is detected. Perform 3 drip applications to the foliage at 7-day intervals.
Almond, Pistachio	Navel Orangeworm (<i>Amyelois transitella</i>)	80	100	Make 3 applications to foliage at 7 to 14-day intervals.
Pome Fruit: Apple, Pear (SL)	Aphid (<i>Aphis pomi</i>) Two-Spotted Mite (<i>Tetranychus urticae</i>) Thrips (<i>Frankliniella occidentalis</i>) Psylla (<i>Cacopsylla bidens</i>) Whitefly (<i>Bemisia tabaci</i>)	32 - 64	80 - 100	Make 3 applications to foliage at 7-days intervals. Use higher rate to higher infestation and when foliage is dense.
Stone Fruit: Apricot, Cherry, Nectarine, Peach, Plum (SL)	Aphid (<i>Aphis pomi</i>) Two-Spotted Mite (<i>Tetranychus urticae</i>) Thrips (<i>Frankliniella occidentalis</i>) Psylla (<i>Cacopsylla bidens</i>) Whitefly (<i>Bemisia tabaci</i>)	32 - 64	80 - 100	Make 3 applications to foliage at 7-days intervals. Use higher rate to higher infestation and when foliage is dense.

All ingredients in this product are exempt from the requirement of a tolerance under 40 CFR 180. () Preharvest Interval (PHI), days between the last application and harvest; NL = No Limit. Re-Entry Intervals (REI): Four (4) hours after application. Its components act as an insecticide and a repellent against insects.