## FIFRA Sec. 2(ee) Recommendation



Reduced use rate for early season control of stripe rust in wheat and triticale grown in Arkansas, Colorado, Idaho, Kansas, Kentucky, Michigan, Minnesota, Mississippi, Missouri, Montana, Nebraska, North Carolina, North Dakota, Ohio, Oklahoma, Oregon, South Dakota, Tennessee, Washington, Wisconsin, and Wyoming.

This Technical Information Bulletin expires on December 31, 2020.

EPA Reg. No. 7969-380

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA.

All applicable directions, restrictions, precautions and Conditions of Sale and Warranty on the EPA registered label are to be followed. Refer to the container label for additional instructions. Always read and follow label directions.

Information contained in this Technical Information Bulletin is not intended to replace or amend any product labeling. Always read and follow all label directions when using any pesticide alone or in tank mix combinations.

The user must have this recommendation in their possession at the time of use.

## **Directions For Use**

Use 3.5 to 7 fl ozs per acre for control of stripe rust (*Puccinia striiformis* f. sp. *tritici*). Follow all use directions under the wheat and triticale section of the **Nexicor™ Xemium® Brand Fungicide** container label.

**Xemium** is a registered trademark of BASF. **Nexicor** is a trademark of BASF.

© 2017 BASF Corporation All rights reserved.

007969-00380.20120170508.NVA 2017-04-523-0083

