# Product Bulletin



**Dow AgroSciences LLC** 

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Closer SC®

EPA Reg. No. 62719-623

## Special 2(ee) Recommendation<sup>†</sup>

(For Distribution and Use in the State of Idaho, Michigan, Oregon, Pennsylvania, Utah and Washington)

## For Control of Mealybugs in Pome (Crop Group 11) and Stone Fruits (Crop Group 12)

Refer to product label for Closer SC for general use precautions, mixing directions, application directions, and restrictions.

#### **ATTENTION**

- <sup>†</sup>This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.
- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Closer SC insecticide before applying. Carefully follow all
  precautionary statements and applicable use directions except as specified below.
- Use of Closer SC according to this Section 2(ee) recommendation is subject to all use precautions and limitations imposed by the label affixed to the container for Closer SC insecticide.

### **Directions for Use**

Pests	Closer SC (fl oz/acre)
mealybugs	4.0 - 5.75
-	(0.063 - 0.09 lb ai/acre)

Apply 4.0 - 5.75 fl oz of Closer SC per acre to provide control of mealybugs in pome fruits. For heavier infestations, use a higher rate in the rate range. Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional local use recommendations for your area.

®Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

R390-020 Issued: 2/24/17 Initial printing