



We create chemistry

SPECIMEN

Group 44 Fungicide

Serifel®

Biofungicide

Serifel is an agricultural biofungicide/bactericide product for suppression of plant diseases in listed crops.



Active Ingredient:

Bacillus amyloliquefaciens strain MBI 600*† 11.0%

Other Ingredients: 89.0%

Total: 100.0%

*Serifel contains a minimum of 5.5×10^{10} colony forming units per gram.

†Formerly named *Bacillus subtilis* strain MBI 600

EPA Reg. No. 71840-18

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN
CAUTION/PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See inside for complete **First Aid, Precautionary Statements, Directions For Use, Conditions of Sale and Warranty**, and state-specific crop and/or use site restrictions.

In case of an emergency endangering life or property involving this product, call day or night 1-800-832-HELP (4357).

Batch Code: (Printed on Bottle)

Net Weight:

BASF Corporation
26 Davis Drive, Research Triangle Park, NC 27709

FIRST AID	
If in eyes	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOTLINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact BASF Corporation for emergency medical treatment information: 1-800-832-HELP (4357).	

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear protective eyewear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Protective eyewear
- A NIOSH-approved particulate respirator with any N, R, or P filter with NIOSH approval number prefix TC-84A; or a NIOSH-approved powered air purifying respirator with an HE filter with NIOSH approval number prefix TC-21C. (Repeated exposure to high concentrations of microbial proteins can cause allergic sensitization.)

Follow the manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

In Case of Emergency

In case of large-scale spill of this product, call:

- CHEMTREC 1-800-424-9300
- BASF Corporation 1-800-832-HELP (4357)

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment
- Your local poison control center (hospital)
- BASF Corporation 1-800-832-HELP (4357)

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval (REI). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **4 hours**.

Exception: If the product is soil injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water) is:

- Coveralls
- Chemical resistant gloves (made of any waterproof material)
- Shoes plus socks
- Protective eyewear

Product Information

This package contains **Serifel**®, a wettable powder (WP). The active ingredient in **Serifel** is *Bacillus amyloliquefaciens* strain MBI 600, a beneficial microbe providing suppression of listed foliar and soilborne plant diseases. Apply **Serifel** preventively in low to medium disease pressure situations. **Serifel** can be applied as a foliar spray, a directed or banded spray, a soil or plant drench, or as an in-furrow application. **Serifel** can be applied alone or in a tank mix with other registered pest control products, unless otherwise noted in the **Crop-Specific Directions**. Apply **Serifel** in a regularly scheduled protective spray program and use in a rotation program with other registered pesticide products. Under higher disease pressure situations, **Serifel** can be tank-mixed with other registered pesticide products.

Label statement required by the State of Oregon

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.html>

Integrated Pest Management (IPM)

Serifel can be integrated into an overall disease and pest management program. Follow cultural practices known to reduce disease development. Consult your local extension specialist, certified crop advisor and/or BASF representative for additional IPM strategies established for your area. **Serifel** may be used in agricultural extension advisory (disease forecasting) programs, which recommend application timing based on environmental factors favorable for disease development.

Application Instructions

Apply rates of **Serifel** as instructed by the **Crop-Specific Directions**. Apply **Serifel** with ground equipment. Equipment must be checked frequently for calibration. If heavy rainfall or irrigation occurs shortly after application, reapplication of **Serifel** may be necessary.

Under low-level disease conditions, the minimum application rates can be used while maximum application rates and shortened spray schedules are to be used for severe or threatening disease conditions. Tank mixing **Serifel** with other registered pesticide products can also improve performance.

Cleaning Spray Equipment

Spraying equipment must be cleaned thoroughly before and after applying this product, particularly if a product with potential to injure crops was used prior to **Serifel**.

Consult BASF Representatives for additional information

regarding agitation and recirculation.

Foliar Ground Applications

Apply **Serifel** in sufficient water to ensure thorough coverage of foliage, bloom, and fruit. Thorough coverage is required. Complete coverage of the stem, all the way down to the soil, is required for suppression of soilborne diseases of the stem.

Maintain agitation of product during the application process. Apply the product mixture shortly after mixing. **DO NOT** store mixed slurries of **Serifel** overnight.

In-Furrow, Shank-In, Injected or Soil Drench Applications

Apply **Serifel** as a water-based suspension alone or with other registered pest control products and fertilizers labeled for in-furrow or soil drench via standard agricultural application machinery. Prior to mixing, determine physical compatibility by mixing proportional quantities of the products in water as described in the **Compatibility Test for Tank Mix Components** section of this label. Maintain agitation of product during the application process. Apply the product mixture shortly after mixing. **DO NOT** store mixed slurries of **Serifel** overnight.

For in-furrow, shanked-in, injected, or soil drench applications, mix 4-16 ounces by weight of **Serifel** with a water volume appropriate for the crop and application type. Use higher rates and consider mixing with other registered pesticide products when conditions favor heavy disease development.

For soil drench applications at planting for seeding or transplants, apply a spray mixture of **Serifel** with adequate water volume to thoroughly drench through the root zone.

For shanked-in and injected applications, the use of **Serifel** can be prior to planting, at planting or after planting of seed or transplants.

For in-furrow applications, apply **Serifel** as an in-furrow spray using the appropriate amount of water for the specific crop. Making a pre-slurry suspension of **Serifel** may help disperse **Serifel** and improve equipment compatibility at lower application volumes.

Instructions for In-Furrow, Shank-In, Injected or Soil Drench Applications of Serifel®										
Rate Per 1000 row feet	Serifel Rate (oz/A)									
(oz product)	12-inch rows	15-inch rows	20-inch rows	22-inch rows	30-inch rows	32-inch rows	34-inch rows	36-inch rows	38-inch rows	40-inch rows
0.3	13.1	10.4	7.8	7.1	5.2	4.9	4.6	4.4	4.1	4.0
0.4	See footnote ¹	13.9	10.4	9.5	7.0	6.5	6.1	5.8	5.5	5.2
0.5	See footnote ¹	See footnote ¹	13.1	11.9	8.7	8.2	7.7	7.3	6.9	6.6
0.6	See footnote ¹	See footnote ¹	15.7	14.2	10.4	9.8	9.2	8.7	8.2	7.9
0.7	See footnote ¹	See footnote ¹	See footnote ¹	See footnote ¹	12.2	11.4	10.7	10.2	9.6	9.2
0.8	See footnote ¹	See footnote ¹	See footnote ¹	See footnote ¹	13.9	13.0	12.3	11.6	11.0	10.5
0.9	See footnote ¹	See footnote ¹	See footnote ¹	See footnote ¹	15.7	14.7	13.8	13.1	12.3	11.8
1.0	See footnote ¹	See footnote ¹	See footnote ¹	See footnote ¹	See footnote ¹	See footnote ¹	15.4	14.5	13.7	13.1
1.1	See footnote ¹	See footnote ¹	See footnote ¹	See footnote ¹	See footnote ¹	See footnote ¹	See footnote ¹	16.0	15.1	14.4
1.2	See footnote ¹	See footnote ¹	See footnote ¹	See footnote ¹	See footnote ¹	See footnote ¹	See footnote ¹	See footnote ¹	See footnote ¹	15.7
Application Directions. Use 0.3 to 1.2 oz of Serifel per 1000 feet of row. Refer to this chart to determine the rate per acre. Apply at planting as an in-furrow application by directing the spray into the furrow before seed is covered.										
DO NOT apply more than 16 oz (1 lb) per acre of Serifel .										
¹ For 40-inch rows, use a maximum of 1.2 oz of Serifel per 1000 row feet. For 36- to 38-inch rows, use a maximum of 1.1 oz of Serifel per 1000 row feet. For 34-inch rows, use a maximum of 1.0 oz of Serifel per 1000 row feet. For 30- to 32-inch rows, use a maximum of 0.9 oz of Serifel per 1000 row feet. For 20- to 22-inch rows, use a maximum of 0.6 oz of Serifel per 1000 row feet. For 15-inch rows, use a maximum of 0.4 oz of Serifel per 1000 row feet. For 12-inch rows, use a maximum of 0.3 oz of Serifel per 1000 row feet.										

Instructions for Directed or Banded Sprays Related to Ground Applications

The application rates shown in the **Crop-Specific Directions** on this label reflect the amount of product to be applied uniformly over an acre of ground on a broadcast basis. In some crops, **Serifel**® may be used as a directed or banded spray over the rows or plant beds with the alleys or row middles left unsprayed. For such uses, reduce the labeled **Serifel** rates in proportion to the area actually sprayed. This adjustment is necessary to avoid applying the product at use rates higher than permitted according to label directions. The following formula may be used to determine the broadcast equivalent rate for doing directed or banded sprays:

sprayed bed width + unsprayed row middle width = total row width

$$\frac{\text{sprayed bed width in inches}}{\text{total row width in inches}} \times \frac{\text{broadcast rate}}{\text{treated acre}} = \frac{\text{band rate}}{\text{field acre}}$$

Example: A directed spray application will be made to 45-inch plant beds that are separated by 15-inch unsprayed row middles.

45 inches sprayed bed width + 15 inches unsprayed row middles = 60 inches total row width

The calculation to determine the appropriate equivalent rate of product to use for this situation based on a label broadcast rate of 4 ounces/acre follows:

$$\frac{\text{45 inch sprayed bed width}}{\text{60 inch total row width}} \times \frac{\text{4 oz Serifel}}{\text{treated acre}} = \frac{\text{3 oz Serifel}}{\text{field acre}}$$

Directions for Use through Sprinkler Chemigation Systems

Sprayer Preparation

Clean chemical tank and injector system thoroughly. Flush system with clean water.

Application Instructions

Apply **Serifel** at rates and timings as described in this label.

Use Precautions for Sprinkler Irrigation Applications

- Apply this product only through sprinkler (including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move) irrigation systems. **DO NOT** apply this product through any other type of irrigation system.
- Add **Serifel** to the pesticide supply tank under agitation containing sufficient water to maintain a continuous flow by the injection equipment. In continuous moving systems, inject this product-water mixture continuously, applying the labeled rate per acre for that crop.
- **DO NOT** exceed 1/2 inch (13,577 gallons) per acre. In stationary or non-continuous moving systems, inject the product-water mixture in the last 15 to 30 minutes of each set allowing sufficient time for all of the required pesticide to be applied by all the sprinkler heads and applying the labeled rate per acre for that crop.
- **DO NOT** apply when wind speed favors drift beyond the area intended for treatment.
- Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. Thorough coverage of foliage is required for good suppression. Maintain good agitation during the entire application period.
- If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.
- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water.
- **DO NOT** connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Specific Instructions for Public Water Systems

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. The pesticide supply tank must be under agitation containing sufficient water to maintain a continuous flow by the injection equipment. In continuous moving systems, inject this product-water mixture continuously, applying the labeled rate per acre for that crop.
8. **DO NOT** apply when wind speed favors drift beyond the area intended for treatment.

Additives and Tank Mixing Information

Serifel® can be tank mixed with most recommended registered pest control products, liquid fertilizers, biological control products, adjuvants, and additives as specified in **Crop-Specific Directions**. Before using any tank mix (bactericides, fungicides, insecticides, herbicides, liquid fertilizers, biological control products, adjuvants, and additives), test the combination on a small portion of the crop to be treated to ensure that a phytotoxic response will not occur as a result of application.

Consult a BASF representative or local agricultural authorities for more information concerning additives.

If tank mixtures are used, adhere to restrictions due to rates and precautions on all labels.

Compatibility Test for Tank Mix Components

Using a suitable container, add proportional amounts of product to water following the tank mixing order below.

TANK MIX COMPONENTS

1. **Water.** For 100 gallons per acre spray volume, use 16 cups (1 gallon) of water. For other spray volumes, adjust rates accordingly. Use only water from the intended source at the source temperature.
2. **Water-dispersible products** (dry flowables, wettable powders, suspension concentrates, or suspoemulsions). Cap the jar and invert 10 cycles.
3. **Water-soluble products.** Cap the jar and invert 10 cycles.
4. **Emulsifiable concentrates** (oil concentrate or methylated seed oil when applicable). Cap the jar and invert 10 cycles.
5. **Water-soluble additives.** Cap the jar and invert 10 cycles.
6. Let the solution stand for 5 minutes.

Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. Simulate tank agitation by inverting the jar for another 10 cycles to ensure that the mixture resuspends. If the tank mix combination does not settle or can be resuspended by agitation, it is considered physically compatible. **DO NOT** use any spray solution that could clog spray nozzles.

Mixing Order

1. **Water.** Begin by agitating a thoroughly clean sprayer tank 3/4 full of clean water. The pH of the spray solution must be between 4 and 9. The product mixture must be applied shortly after mixing. **DO NOT** store mixed slurries of **Serifel** overnight.
2. **Agitation.** Maintain constant agitation throughout mixing and application.
3. **Inductor.** If an inductor is used, rinse it thoroughly after each component has been added.
4. **Products in polyvinyl alcohol (PVA) bags.** Place any product contained in water-soluble PVA bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
5. **Water-dispersible products** (such as **Serifel**, dry flowables, wettable powders, suspension concentrates, or suspoemulsions).
6. **Water-soluble products.**
7. **Emulsifiable concentrates** (such as oil concentrates when applicable).
8. **Water-soluble additives** (such as ammonium sulfate (AMS) or urea ammonium nitrate (UAN) when

applicable).

9. **Remaining quantity of water.**

Make sure that each component is thoroughly mixed and suspended before adding tank mix partners. Maintain constant agitation during application. See section **Crop-Specific Directions** for more details.

Restrictions and Limitations

- **Crop Rotation Restriction** - None
- **Preharvest Interval (PHI)** – 0 Day
- **Re-entry Interval (REI)** – 4 hours
- **Not for use** in greenhouse or transplant production
- **Not for use** in California on crops marked with an asterisk (*) in **Crop-Specific Directions**

Crop-Specific Directions

Crop	Target Diseases	Product Use Rate per Application (oz or lb of Serifel®/Acre)	Application Directions
Alfalfa and Clover Includes alfalfa and clover mixed with forage grasses	Foliar diseases White mold (<i>Sclerotinia sclerotiorum</i>)	4 – 16 oz/A (0.25 – 1.0 lb/A)	White mold: Begin application shortly after emergence or transplanting and repeat on 7-10 day intervals as needed. Mix and apply Serifel in sufficient water volume to ensure uniform dispersion in spray tank and thorough coverage of foliage and shoot tissue.

Crop-Specific Directions (*continued*)

Crop	Target Diseases	Product Use Rate per Application (oz or lb of Serifel®/Acre)	Application Directions
Artichoke	Foliar diseases Anthracnose <i>(Colletotrichum spp.)</i> Alternaria leaf spot <i>(Alternaria spp.)</i> Botrytis gray mold <i>(Botrytis spp.)</i> Powdery mildew <i>(Leveillula taurica)</i> Ramularia leaf spot* <i>(Ramularia cynaræ)</i>	4 – 16 oz/A (0.25 – 1.0 lb /A)	Begin foliar applications before infection and continue on 7-10 day intervals if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high. Mix and apply Serifel in sufficient water volume to ensure uniform dispersion in spray tank and thorough coverage of foliage and shoot tissue.
	Soil diseases Fusarium wilt <i>(Fusarium spp.)</i> Pythium damping off <i>(Pythium spp.)</i> Rhizoctonia crown rot <i>(Rhizoctonia solani)</i> Verticillium wilt <i>(Verticillium spp.)</i>	4 – 16 oz/A (0.25 – 1.0 lb /A)	See Application Instructions for In-Furrow, Drench, Shank-In and Injected Applications. Apply a high enough water volume to thoroughly soak soil through the root zone.

*Not registered for use in California.

Crop-Specific Directions (*continued*)

Crop	Target Diseases	Product Use Rate per Application (oz or lb of Serifel®/Acre)	Application Directions
Asparagus	Foliar diseases Anthracnose <i>(Colletotrichum spp.)</i> Alternaria leaf spot <i>(Alternaria spp.)</i> Botrytis gray mold <i>(Botrytis spp.)</i> Phytophthora spear and crown rot <i>(Phytophthora spp.)</i> Watery soft rot <i>(Sclerotinia spp.)</i>	4 – 16 oz/A (0.25 – 1.0 lb/A)	Begin foliar applications prior to infection and continue on 7-10 day intervals if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high. Mix and apply Serifel in sufficient water volume to ensure uniform dispersion in spray tank and thorough coverage of foliage and shoot tissue.
	Soil diseases Fusarium wilt <i>(Fusarium spp.)</i> Pythium damping off <i>(Pythium spp.)</i> Rhizoctonia root rot <i>(Rhizoctonia solani)</i>	4 – 16 oz/A (0.25 – 1.0 lb/A)	See Application Instructions for In-Furrow, Drench, Shank-In and Injected Applications. Apply a high enough water volume to thoroughly soak soil through the root zone.

Crop-Specific Directions (*continued*)

Crop	Target Diseases	Product Use Rate per Application (oz or lb of Serifel®/Acre)	Application Directions
Berries and small fruits subgroups Bushberry subgroup Blueberry (highbush and lowbush) Currant Elderberry Gooseberry Huckleberry Caneberry subgroup Blackberry (all varieties) Loganberry Raspberry (black and red) Wild raspberry Low growing berry subgroup (except strawberry) Bearberry Bilberry Cloudberry Cranberry Lingonberry Muntries Partridgeberry Small fruit vine climbing subgroup (except fuzzy kiwifruit and grape) Amur river grape Gooseberry Kiwifruit, hardy Maypop Schisandra berry	Foliar diseases Alternaria fruit rot <i>(Alternaria spp.)</i> Anthracnose <i>(Colletotrichum spp.)</i> Bacterial canker <i>(Pseudomonas spp.)</i> Botrytis gray mold <i>(Botrytis spp.)</i> Mummy berry <i>(Monilinia spp.)</i> Powdery mildew <i>(Sphaerotheca spp., Microsphaera spp., Podosphaera spp.)</i>	4 – 16 oz/A (0.25 – 1.0 lb/A)	For Mummy berry, begin applications at bud break and continue on a 7-10 day interval as needed. For Alternaria fruit rot, Anthracnose, Botrytis gray mold, and Powdery mildew, begin applications prior to infection and continue on 2 - 10 day intervals if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high. For Bacterial canker, apply before fall rains and again during dormancy prior to spring growth. Apply prior to disease development and continue on 2 – 10 day intervals as needed throughout the growing season. For Cranberries, make application only to non-flooded fields. Mix and apply Serifel in sufficient water volume to ensure uniform dispersion in spray tank and thorough coverage of foliage and shoot tissue.

Crop-Specific Directions (*continued*)

Crop	Target Diseases	Product Use Rate per Application (oz or lb of Serifel®/Acre)	Application Directions
Brassica (cole) leafy vegetables group Head and stem Broccoli Broccoli, Chinese Brussel sprouts Cabbage Cabbage, Chinese Cabbage, Chinese mustard Cauliflower Cavalo broccolo Kohlrabi Leafy greens Broccoli raab Chinese cabbage (bok choy) Collards Kale Mizuna Mustard greens Mustard spinach Rape greens	Foliar diseases Alternaria leaf spot <i>(Alternaria spp.)</i> Botrytis gray mold <i>(Botrytis spp.)</i> Downy mildew <i>(Peronospora spp.)</i> Powdery mildew <i>(Erysiphe polygoni)</i> Pin rot <i>(Alternaria spp./Xanthomonas spp. complex)</i> Rust <i>(Puccinia porri)</i> White rust* <i>(Albugo candida)</i>	4 – 16 oz/A (0.25 – 1.0 lb/A)	Begin foliar applications prior to infection and continue on 3-10 day intervals if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high. Mix and apply Serifel in sufficient water volume to ensure uniform dispersion in spray tank and thorough coverage of foliage and shoot tissue.
	Soil diseases Fusarium wilt <i>(Fusarium spp.)</i> Phytophthora root rot <i>(Phytophthora spp.)</i> Pythium damping off <i>(Pythium spp.)</i> Rhizoctonia root rot <i>(Rhizoctonia solani)</i> Verticillium wilt <i>(Verticillium spp.)</i>	4 – 16 oz/A (0.25 – 1.0 lb/A)	See Application Instructions for In-Furrow, Drench, Shank-In and Injected Applications. Apply a high enough water volume to thoroughly soak soil through the root zone.

*Not registered for use in California.

Crop-Specific Directions (continued)

Crop	Target Diseases	Product Use Rate per Application (oz or lb of Serifel®/Acre)	Application Directions
Bulb vegetables group Chive, fresh leaves Chive, Chinese, fresh leaves Daylily, bulb Elegans hosta Fritillaria, bulb Fritillaria, leaves Garlic, bulb Garlic, great-headed, bulb Garlic, serpent, bulb Kurrat Lady's leek Leek Leek, wild Lily, bulb Onion, Beltsville bunching Onion, bulb Onion, Chinese, bulb Onion, fresh Onion, green Onion, macrostem Onion, pearl Onion, potato, bulb Onion, tree, tops Onion, Welsh, tops Shallot, bulb Shallot, fresh, leaves	Foliar diseases Botrytis leaf blight and neck rot (<i>Botrytis</i> spp.) Downy mildew (<i>Peronospora</i> spp.) Purple blotch (<i>Alternaria porri</i>) Powdery mildew (<i>Erysiphe</i> spp.) Stemphylium leaf blight and stalk rot* (<i>Stemphylium vesicarium</i>) White rot (<i>Sclerotium cepivorum</i>)	4 – 16 oz/A (0.25 – 1.0 lb/A)	Begin foliar applications prior to infection and continue on 7-10 day intervals if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high. Mix and apply Serifel in sufficient water volume to ensure uniform dispersion in spray tank and thorough coverage of foliage and shoot tissue.
	Soil diseases Fusarium wilt (<i>Fusarium</i> spp.) Pink root* (<i>Phoma</i> spp.) Pythium damping off (<i>Pythium</i> spp.) Rhizoctonia root rot (<i>Rhizoctonia solani</i>)	4 – 16 oz/A (0.25 – 1.0 lb/A)	See Application Instructions for In-Furrow, Drench, Shank-In and Injected Applications. Apply a high enough water volume to thoroughly soak soil through the root zone.

*Not registered for use in California.

Crop-Specific Directions (continued)

Crop	Target Diseases	Product Use Rate per Application (oz or lb of Serifel®/Acre)	Application Directions
Citrus fruit group Calamondin Citrus citron Citrus hybrids Grapefruit Kumquat Lemon Lime Mandarin Orange (sour and sweet) Pummelo Satsuma Tangelo Tangerine Tangor	Foliar diseases Alternaria leaf spot <i>(Alternaria spp.)</i> Citrus canker <i>(Xanthomonas campestris pv. citri)</i> Greasy spot <i>(Mycosphaerella citri)</i> Melanose <i>(Diaporthe citri)</i> Post bloom fruit drop <i>(Colletotrichum spp.)</i> Scab <i>(Elsinoe fawcetti)</i>	4 – 16 oz/A (0.25 – 1.0 lb/A)	For Greasy spot, begin applications at the start of each new flush of foliage and repeat for each new flush. Tank mix Serifel with labeled spray oil or copper based fungicide products. For Post bloom fruit drop, begin applications at early bloom when conditions are favorable for disease development. Continue on a 7-10 day interval as needed. For Alternaria leaf spot, begin foliar applications prior to infection and continue on 7-10 day intervals if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high. For Melanose, begin applications at petal fall and continue on a 14-21 day interval as needed. For Citrus canker, apply when environmental conditions are conducive to disease development and continue on a 7 – 14 day interval if needed. For Scab, apply at the first new flush of foliage and repeat at petal fall and when fruit are ½ inch in diameter. Mix and apply Serifel in sufficient water volume to ensure uniform dispersion in spray tank and thorough coverage of foliage and shoot tissue.
	Soil diseases Phytophthora root rot <i>(Phytophthora spp.)</i> Pythium damping off <i>(Pythium spp.)</i>	4 – 16 oz/A (0.25 – 1.0 lb/A)	See Application Instructions for In-Furrow, Drench, Shank-In and Injected Applications. Apply a high enough water volume to thoroughly soak soil through the root zone.

Crop-Specific Directions (continued)

Crop	Target Diseases	Product Use Rate per Application (oz or lb of Serifel®/Acre)	Application Directions
Cucurbit vegetables group Chayote Chinese waxgourd Citron melon Cucumber Gherkin Pumpkin Watermelon Edible gourd Hyotan Cucuzza Chinese okra Momordica spp. Balsam apple Balsam pear Bitter melon Chinese cucumber Muskmelon Cantaloupe Casaba Crenshaw melon Golden pershaw melon Honeydew melon Honey balls Mango melon Persioan melon Pineapple melon Santa Claus melon Snake melon Summer squash Crookneck squash Scallop squash Straightneck squash Vegetable marrow Zucchini Winter squash Butternut squash Calabaza Hubbard squash Acorn squash Spaghetti squash	Foliar diseases Anthracnose <i>(Colletotrichum spp.)</i> Alternaria leaf spot <i>(Alternaria spp.)</i> Downy mildew <i>(Pseudoperonospora spp.)</i> Gummy stem blight <i>(Didymella bryoniae)</i> Powdery mildew <i>(Erysiphe spp., Sphaerotheca spp.)</i>	4 – 16 oz/A (0.25 – 1.0 lb/A)	Begin foliar applications prior to infection and continue on 7-10 day intervals if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high. Mix and apply Serifel in sufficient water volume to ensure uniform dispersion in spray tank and thorough coverage of foliage and shoot tissue.
	Soil diseases Charcoal rot* <i>(Macrophomina phaseolina)</i> Fusarium wilt <i>(Fusarium spp.)</i> Phytophthora root rot <i>(Phytophthora spp.)</i> Pythium damping off <i>(Pythium spp.)</i> Rhizoctonia root rot <i>(Rhizoctonia solani)</i> Verticillium wilt <i>(Verticillium spp.)</i> Vine decline* <i>(Monosporascus cannonballus)</i>	4 – 16 oz/A (0.25 – 1.0 lb/A)	See Application Instructions for In-Furrow, Drench, Shank-In and Injected Applications. Apply a high enough water volume to thoroughly soak soil through the root zone.

*Not registered for use in California.

Crop-Specific Directions (continued)

Crop	Target Diseases	Product Use Rate per Application (oz or lb of Serifel®/Acre)	Application Directions
Corn Field corn Pop corn Silage corn Sweet corn Seed production	Foliar diseases Common rust (<i>Puccinia sorghi</i>) Southern leaf blight (<i>Bipolaris maydis</i>)	4 – 16 oz/A (0.25 – 1.0 lb/A)	Begin foliar applications prior to infection and continue on 7-10 day intervals if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high. Mix Serifel with a labeled corn fungicide. Mix and apply Serifel in sufficient water volume to ensure uniform dispersion in spray tank and thorough coverage of foliage and shoot tissue.
	Soil diseases Charcoal rot (<i>Macrophomina phaseolina</i>) Fusarium wilt (<i>Fusarium</i> spp.) Phytophthora root rot (<i>Phytophthora</i> spp.) Pythium damping off (<i>Pythium</i> spp.) Rhizoctonia root rot (<i>Rhizoctonia solani</i>)	4 – 16 oz/A (0.25 – 1.0 lb/A)	See Application Instructions for In-Furrow, Drench, Shank-In and Injected Applications. Apply a high enough water volume to thoroughly soak soil through the root zone.

*Not registered for use in California.

Crop-Specific Directions (continued)

Crop	Target Diseases	Product Use Rate per Application (oz or lb of Serifel®/Acre)	Application Directions
Fruiting vegetables group Eggplant Groundcherry Pepino Pepper (all varieties) Tomatillo Tomato	Foliar diseases Anthracnose (<i>Colletotrichum</i> spp.) Botrytis gray mold (<i>Botrytis</i> spp.) Bacterial speck (<i>Pseudomonas</i> spp.) Bacterial spot (<i>Xanthomonas</i> spp.) Buck-eye rot (<i>Phytophthora parasitica</i>) Early blight (<i>Alternaria</i> spp.) Late blight (<i>Phytophthora infestans</i>) Powdery mildew (<i>Leveillula</i> spp., <i>Oidiopsis</i> spp., <i>Erysiphe</i> spp., <i>Sphaerotheca</i> spp.) Target spot* (<i>Corynespora cassiicola</i>)	4 – 16 oz/A (0.25 – 1.0 lb/A)	For Bacterial spot, Bacterial speck and Target spot, begin applications shortly after emergence or transplanting and continue on a 2 – 7 day application interval if conditions are conducive to disease development. For improved suppression of bacterial spot and speck, tank mix or rotate with labeled copper based bactericides. For Anthracnose, Botrytis gray mold, Buck-eye rot and Powdery mildew, begin application shortly after emergence or transplanting and continue on 7-10 day intervals as needed. For Early blight and Late blight, apply when plants are 4-6 inches tall and continue on 5-10 day intervals as needed. Mix and apply Serifel in sufficient water volume to ensure uniform dispersion in spray tank and thorough coverage of foliage and shoot tissue.
	Soil diseases Fusarium wilt (<i>Fusarium</i> spp.) Phytophthora root rot (<i>Phytophthora</i> spp.) Pythium damping off (<i>Pythium</i> spp.) Rhizoctonia root rot (<i>Rhizoctonia solani</i>) Sclerotinia stem rot* (<i>Sclerotinia sclerotiorum</i>) Southern blight (<i>Sclerotium rolfsii</i>) Verticillium wilt (<i>Verticillium</i> spp.)	4 – 16 oz/A (0.25 – 1.0 lb/A)	See Soil Application Instructions for In-Furrow, Drench, Shank-In and Injected Applications. Apply a high enough water volume to thoroughly soak soil through the root zone. Complete coverage of the stem, all the way down to the soil, is required for suppression of soilborne diseases of the stem.

*Not registered for use in California.

Crop-Specific Directions (continued)

Crop	Target Diseases	Product Use Rate per Application (oz or lb of Serifel®/Acre)	Application Directions
Grape	Foliar diseases Botryosphaeria dieback, including Black dead arm, Botryosphaeria canker and Diplodia cane dieback* (<i>Botryosphaeria</i> spp., <i>Diplodia</i> spp., <i>Lasiodiplodia</i> spp., <i>Neofusicoccum</i> spp., <i>Dothiorella</i> spp. and <i>Sphaeropsis</i> spp.) Botrytis gray mold (<i>Botrytis</i> spp.) Downy mildew (<i>Plasmopara viticola</i>) Esca, Black measles* (<i>Phaeoacremonium</i> spp., <i>Phaeomoniella</i> spp.) Eutypa (<i>Eutypa lata</i>) Macrophoma rot* (<i>Botryosphaeria dothidea</i>) Phomopsis cane and leaf spot (<i>Phomopsis viticola</i>) Phomopsis dieback (<i>Phomopsis viticola</i>) Phomopsis (<i>Phomopsis viticola</i>) Powdery mildew (<i>Uncinula</i> spp., <i>Erysiphe</i> spp.) Summer bunch rot (<i>Aspergillus</i> spp.*, <i>Alternaria</i> spp., <i>Botrytis</i> spp., <i>Cladosporium</i> spp., <i>Penicillium</i> spp.*, <i>Rhizopus</i> spp.)	4 – 16 oz/A (0.25 – 1.0 lb/A)	For Powdery mildew and Phomopsis, begin foliar applications at bud break or prior to onset of disease. Repeat applications on 5 –10 day intervals as needed. For Downy mildew, begin applications before pre-bloom and continue on 7–10 day intervals as needed. For Botrytis and Summer bunch rot, begin applications prior to disease development when conditions favor the development of Botrytis and summer bunch rot such as during early bloom, bunch preclosure and veraison. Use higher rates and shorter intervals when disease pressure is high. For Botryosphaeria dieback, Phomopsis dieback, Esca and Eutypa, apply to pruning wounds. Sanitation of pruning tools is critical. For Macrophoma rot, apply to pruning wounds. Sanitation of pruning tools is critical. For foliar symptom suppression during the growing season, apply just after bloom and continue on 7 – 10 day intervals until harvest when environmental conditions are conducive for disease development. For Phomopsis cane and leaf spot, begin applications between bud break and 1 inch shoots and repeat when shoots are 6-8 inches long. Mix and apply Serifel in sufficient water volume to ensure uniform dispersion in spray tank and thorough coverage of foliage and shoot tissue.

*Not registered for use in California.

Crop-Specific Directions (continued)

Crop	Target Diseases	Product Use Rate per Application (oz or lb of Serifel®/Acre)	Application Directions
Hops	Foliar diseases Powdery mildew <i>(Sphaerotheca macularis)</i>	4 – 16 oz/A (0.25 – 1.0 lb/A)	For Powdery mildew, begin foliar applications prior to onset of disease and continue on 5 –10 day intervals as needed. Mix and apply Serifel in sufficient water volume to ensure uniform dispersion in spray tank and thorough coverage of foliage and shoot tissue.

Crop-Specific Directions (continued)

Crop	Target Diseases	Product Use Rate per Application (oz or lb of Serifel®/Acre)	Application Directions
Leafy vegetables (except <i>Brassica</i> vegetables) group Amaranth Arugula Cardoon Celery Celery, Chinese Celtuce Chervil Chrysanthemum (edible-leaved and garland) Corn salad Cress (garden and upland) Dandelion Dock Endive Fennel, Florence Lettuce (head and leaf) Orach Parsley Purslane (garden and winter) Radicchio (red chicory) Rhubarb Spinach Spinach (New Zealand and vine) Swiss chard	Foliar diseases Bacterial leaf spot (<i>Xanthomonas</i> and <i>Pseudomonas</i> spp.) Botrytis gray mold (<i>Botrytis</i> spp.) Cercospora leaf spot (<i>Cercospora</i> spp.) Downy mildew (<i>Bremia lactucae</i> , <i>Peronospora</i> spp.) Head and leaf drop (<i>Sclerotinia</i> spp.) Pink rot (<i>Sclerotinia sclerotiorum</i>) Powdery mildew (<i>Erysiphe</i> spp.) White mold, Lettuce drop (<i>Sclerotinia sclerotiorum</i>) White rust* (<i>Albugo</i> spp.)	4 – 16 oz/A (0.25 – 1.0 lb/A)	For Head and leaf drop, White mold, and Lettuce drop, apply before emergence as a directed or banded spray 4 – 6 inches wide. Apply again at thinning or cultivation and continue on a 10 – 14 day interval if conditions remain conducive for disease development. For Pink rot, apply before emergence as a directed or banded spray 4 – 6 inches wide. Apply again at thinning or cultivation and continue on a 10 – 14 day interval if conditions remain conducive for disease development. Begin application approximately eight weeks before harvest and repeat on a 14-day interval. For Botrytis gray mold, Cercospora leaf spot, Downy mildew, White rust and Powdery mildew, apply prior to infection and continue on 7-10 day intervals as needed. Use the higher rate and shorter interval when disease pressure is high. For Bacterial leaf spot, begin applications prior to disease onset and continue on 2 – 10 day application intervals if conditions are conducive to disease development. Mix and apply Serifel in sufficient water volume to ensure uniform dispersion in spray tank and thorough coverage of foliage and shoot tissue.
	Soil diseases Bottom rot (<i>Rhizoctonia solani</i>) Head and leaf drop (<i>Sclerotinia</i> spp.) Verticillium wilt (<i>Verticillium</i> spp.)	4 – 16 oz/A (0.25 – 1.0 lb/A)	See Soil Application Instructions for In-furrow, Drench, Shank-In and Injected Applications. Apply a high enough water volume to thoroughly soak soil through the root zone.

*Not registered for use in California.

Crop-Specific Directions (continued)

Crop	Target Diseases	Product Use Rate per Application (oz or lb of Serifel®/Acre)	Application Directions
Legume vegetables (succulent and dried beans and peas, except soybean) group Broad bean Chickpea Guar Lablab bean Lentil Pigeon pea Lupinus spp. Grain lupin Sweet lupin White lupin Phaseolus spp. Field bean Kidney bean Lima bean Navy bean Pinto bean Tepary bean Pisum spp. English pea Field pea Garden pea Green pea Vigna spp. Adzuki bean Blackeyed pea Catjang Cowpea Crowder pea Moth bean Mung bean Rice bean Southern pea Urd bean	Foliar diseases White mold (<i>Sclerotinia sclerotiorum</i>) Botrytis gray mold (<i>Botrytis</i> spp.) Powdery mildew (<i>Microsphaera diffusa</i>)	4 – 16 oz/A (0.25 – 1.0 lb/A)	For White mold and Botrytis gray mold , apply shortly after emergence or transplanting when conditions favor disease development. Repeat on 7–10 day intervals as needed. When disease pressure is high, use Serifel in a rotation program with other registered fungicides. For Powdery mildew , begin foliar applications prior to onset of disease and continue on 5 – 10 day intervals as needed. Mix and apply Serifel in sufficient water volume to ensure uniform dispersion in spray tank and thorough coverage of foliage and shoot tissue.
	Soil diseases Fusarium wilt (<i>Fusarium</i> spp.) Phytophthora root rot (<i>Phytophthora</i> spp.) Pythium damping off (<i>Pythium</i> spp.) Rhizoctonia root rot (<i>Rhizoctonia solani</i>) Verticillium wilt (<i>Verticillium</i> spp.)	4 – 16 oz/A (0.25 – 1.0 lb/A)	See Soil Application Instructions for In-furrow, Drench, Shanked-In and Injected Applications. Apply a high enough water volume to thoroughly soak soil through the root zone.

Crop-Specific Directions (*continued*)

Crop	Target Diseases	Product Use Rate per Application (oz or lb of Serifel®/Acre)	Application Directions
Oilseed group Canola Castor Cotton Flax Rapeseed Safflower Sesame Sunflower	Foliar diseases White mold (<i>Sclerotinia sclerotiorum</i>)	4 – 16 oz/A (0.25 – 1.0 lb/A)	For White mold , apply shortly after emergence when conditions favor disease development. Repeat on 7– 10 day intervals as needed. When disease pressure is high, use higher rates and shorter intervals. Mix and apply Serifel in sufficient water volume to ensure uniform dispersion in spray tank and thorough coverage of foliage and shoot tissue.
	Soil diseases Cylindrocladium black rot* (<i>Cylindrocladium</i> spp.) Fusarium wilt (<i>Fusarium</i> spp.) Phytophthora root rot (<i>Phytophthora</i> spp.) Pythium damping off (<i>Pythium</i> spp.) Rhizoctonia root rot (<i>Rhizoctonia solani</i>) Southern blight (<i>Sclerotium rolfsii</i>) Verticillium wilt (<i>Verticillium</i> spp.)	4 – 16 oz/A (0.25 – 1.0 lb/A)	See Soil Application Instructions for In-furrow, Drench, Shank-In and Injected Applications. Apply a high enough water volume to thoroughly soak soil through the root zone.

*Not registered for use in California.

Crop-Specific Directions (*continued*)

Crop	Target Diseases	Product Use Rate per Application (oz or lb of Serifel®/Acre)	Application Directions
Pome fruit group Apple Crabapple Loquat Mayhaw Oriental pear Pear Quince	Foliar diseases Alternaria blotch (<i>Alternaria mali</i>) Bitter rot (<i>Colletotrichum</i> spp.) Blue mold* (<i>Penicillium</i> spp.) Bot rot* (<i>Botryosphaeria dothidea</i>) Botrytis gray mold (<i>Botrytis</i> spp.) Brooks spot (<i>Mycosphaerella pomi</i>) Bull's eye rot* (<i>Neofabraea</i> spp.) Cedar apple rust* (<i>Gymnosporangium juniper-virginianae</i>) Fire blight (<i>Erwinia amylovora</i>) Flyspeck* (<i>Schizothyrium pomi</i>) Powdery mildew (<i>Podosphaera</i> spp.) Scab (<i>Venturia</i> spp.)	4 – 16 oz/A (0.25 – 1.0 lb/A)	For Bitter rot, Bot rot, Brooks spot, Bull's eye rot, Cedar apple rust, and Flyspeck, begin applications at pre-bloom when conditions are favorable for disease development. Repeat applications on 7-14 day intervals or as needed. For Powdery mildew, begin applications at tight cluster or earlier if conditions are favorable for disease development. Repeat application on 7-10 day intervals through the second cover spray. For Fire blight, begin applications at 1-5% bloom and repeat as needed to protect new blossoms when conditions are favorable for disease development. During rapid bloom development, apply at 2-7 day intervals when conditions are favorable for disease development. Continue application at 7-day intervals after petal fall when conditions are favorable for disease development. For improved performance, use Serifel in a rotational program with a registered antibiotic product. For Scab, begin applications at green tip or when conditions are favorable for disease development. Continue applications on 7-10 day intervals as needed. Use Serifel in a rotational program with fungicides registered for scab control. For Alternaria blotch, Blue mold and Botrytis gray mold, begin applications prior to disease development and continue on 7-10 day intervals as needed. Mix and apply Serifel in sufficient water volume to ensure uniform dispersion in spray tank and thorough coverage of foliage and shoot tissue.

*Not registered for use in California.

Crop-Specific Directions (continued)

Crop	Target Diseases	Product Use Rate per Application (oz or lb of Serifel®/Acre)	Application Directions
Soybean <i>(Glycine max)</i>	Foliar diseases White mold <i>(Sclerotinia sclerotiorum)</i>	4 – 16 oz/A (0.25 – 1.0 lb/A)	Begin foliar applications prior to infection and continue on 7-10 day intervals if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high. Mix Serifel with a labeled soybean fungicide. Mix and apply Serifel in sufficient water volume to ensure uniform dispersion in spray tank and thorough coverage of foliage and shoot tissue.
	Soil diseases Charcoal rot <i>(Macrophomina phaseolina)</i> Fusarium wilt <i>(Fusarium spp.)</i> Phytophthora root rot <i>(Phytophthora spp.)</i> Pythium damping off <i>(Pythium spp.)</i> Rhizoctonia root rot <i>(Rhizoctonia solani)</i>	4 – 16 oz/A (0.25 – 1.0 lb/A)	See Application Instructions for In-Furrow, Drench, Shank-In and Injected Applications. Apply a high enough water volume to thoroughly soak soil through the root zone.

Crop-Specific Directions (continued)

Crop	Target Diseases	Product Use Rate per Application (oz or lb of Serifel®/Acre)	Application Directions
Stone fruit group Apricot Cherry (sweet and tart) Nectarine Peach Plum (all varieties) Plumcot Prune	Foliar diseases Alternaria spot <i>(Alternaria alternata)</i> Anthracnose <i>(Colletotrichum spp.)</i> Bacterial canker <i>(Pseudomonas spp.)</i> Bacterial spot <i>(Xanthomonas arboricola)</i> Blossom blight <i>(Monilinia spp.)</i> Botrytis gray mold <i>(Botrytis spp.)</i> Brown rot of fruit <i>(Monilinia spp.)</i> Powdery mildew <i>(Sphaerotheca spp., Podosphaera spp.)</i> Ripe fruit rot <i>(Monilinia spp., Botrytis spp., Rhizopus spp.)</i> Shot hole <i>(Wilsonomyces carpophilus)</i>	4 – 16 oz/A (0.25 – 1.0 lb/A)	For Anthracnose and Brown rot of fruit, begin applications prior to disease development when conditions are favorable for disease development and continue on 7-10 day intervals as needed. For Bacterial canker, apply after harvest prior to fall rain events. Re-apply in the early spring during dormancy. For Bacterial spot, begin applications at bud break and continue at 7-14 day intervals as needed until harvest. For improved performance, use Serifel in a rotational program with registered antibiotic and copper products. Use higher rates and shorter intervals when disease pressure is high. For Blossom blight, begin applications at early bloom and continue through petal fall at 7-day intervals as needed. For Powdery mildew, begin applications at popcorn stage and continue on 7-day intervals as needed. For all other listed diseases, begin foliar applications prior to infection and continue on 7-10 day intervals if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high. Mix and apply Serifel in sufficient water volume to ensure uniform dispersion in spray tank and thorough coverage of foliage and shoot tissue.

Crop-Specific Directions (continued)

Crop	Target Diseases	Product Use Rate per Application (oz or lb of Serifel®/Acre)	Application Directions
Strawberry	Foliar diseases Angular leaf spot <i>(Xanthomonas fragariae)</i> Anthracnose <i>(Colletotrichum spp.)</i> Botrytis gray mold <i>(Botrytis spp.)</i> Leaf scorch* <i>(Diplocarpon earliana)</i> Leaf spot* <i>(Mycosphaerella fragariae)</i> Powdery mildew <i>(Erysiphe spp., Sphaerotheca spp.)</i>	4 – 16 oz/A (0.25 – 1.0 lb/A)	For Angular leaf spot, Anthracnose, Leaf scorch, and Leaf spot, begin foliar applications prior to infection and continue on 7-10 day intervals if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high. For Botrytis and Powdery mildew, begin applications at or before flowering, continuing on 7-10 day intervals as needed. Use higher rates and shorter intervals when disease pressure is high. Use Serifel in a rotational program or tank mix with other registered fungicides labeled for Botrytis and Powdery mildew control. Mix and apply Serifel in sufficient water volume to ensure uniform dispersion in spray tank and thorough coverage of foliage and shoot tissue.
	Soil diseases Charcoal rot <i>(Macrophomina phaseolina)</i> Fusarium wilt <i>(Fusarium spp.)</i> Phytophthora root rot <i>(Phytophthora spp.)</i> Pythium damping off <i>(Pythium spp.)</i> Rhizoctonia root rot <i>(Rhizoctonia solani)</i> Verticillium wilt <i>(Verticillium spp.)</i>	4 – 16 oz/A (0.25 – 1.0 lb/A)	See Soil Application Instructions for In-furrow, Drench, Shank-In and Injected Applications. Apply a high enough water volume to thoroughly soak soil through the root zone.

*Not registered for use in California.

Crop-Specific Directions (continued)

Crop	Target Diseases	Product Use Rate per Application (oz or lb of Serifel®/Acre)	Application Directions
Sugar beet	Foliar diseases Leaf spot <i>(Cercospora beticola)</i> Powdery mildew <i>(Erysiphe betae)</i> Rhizoctonia stem canker and crown rot <i>(Rhizoctonia solani)</i>	4 – 16 oz/A (0.25 – 1.0 lb/A)	For Leaf spot and Powdery mildew, begin applications prior to infection and continue on 7-14 day intervals if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high. For Rhizoctonia stem canker and crown rot, begin applications at the 2 leaf stage and continue on a 3 – 10 day interval until the 8 leaf stage. Mix and apply Serifel in sufficient water volume to ensure uniform dispersion in spray tank and thorough coverage of foliage and shoot tissue.

Crop-Specific Directions (*continued*)

Crop	Target Diseases	Product Use Rate per Application (oz or lb of Serifel®/Acre)	Application Directions
Tobacco	Foliar diseases Angular leaf spot <i>(Pseudomonas spp.)</i> Anthracnose <i>(Colletotrichum destrutivum, Glomerella glycines)</i> Barnspot/Frogeye leaf spot <i>(Cercospora nicotianae)</i> Brown spot <i>(Alternaria alternata)</i> Blue mold <i>(Peronospora spp.)</i> Botrytis gray mold <i>(Botrytis cinerea)</i> Collar rot <i>(Sclerotinia sclerotiorum)</i> Powdery mildew <i>(Erysiphe cichoracearum)</i> Target spot <i>(Rhizoctonia solani)</i>	4 – 16 oz/A (0.25 – 1.0 lb/A)	Begin foliar applications prior to infection and continue on 7-10 day intervals if conditions are conducive for disease development. Use the higher rate and shorter interval when disease pressure is high. Mix and apply Serifel in sufficient water volume to ensure uniform dispersion in spray tank and thorough coverage of foliage and shoot tissue.
	Soil diseases Black root rot <i>(Thielaviopsis basicola)</i> Black shank <i>(Phytophthora spp.)</i> Charcoal rot <i>(Macrophomina phaseolina)</i> Southern blight <i>(Sclerotium rolfsii)</i>	4 – 16 oz/A (0.25 – 1.0 lb/A)	See Soil Application Instructions for In-furrow, Drench, Shank-In and Injected Applications. Apply in enough water to thoroughly soak soil through the root zone.

Crop-Specific Directions (continued)

Crop	Target Diseases	Product Use Rate per Application (oz or lb of Serifel®/Acre)	Application Directions
Root and tuber vegetables (except sugar beet) group Potato Arracacha Arrowroot Carrot Cassava (bitter and sweet) Chayote Chinese artichoke Chufa Dasheen (taro) Edible canna Ginger Jerusalem artichoke Leren Sweet potato Tanier True yam Turmeric Yam bean	Foliar diseases Aerial stem rot* <i>(Erwinia carotovora)</i> Bacterial leaf spot <i>(Xanthomonas campestris pv. carotae)</i> Black dot <i>(Colletotrichum coccodes)</i> Black rot/Crown rot <i>(Alternaria spp.)</i> Brown spot and black pit <i>(Alternaria alternata)</i> Early blight <i>(Alternaria solani)</i> Botrytis gray mold <i>(Botrytis spp.)</i> Late blight <i>(Phytophthora infestans)</i> White mold <i>(Sclerotinia sclerotiorum)</i>	4 – 16 oz/A (0.25 – 1.0 lb/A)	For Black rot/Crown rot, Bacterial leaf spot and Botrytis gray mold, begin foliar applications shortly after emergence or transplanting and continue on 7-10 day intervals if conditions are favorable for disease development. Use the higher rate and shorter interval when disease pressure is high. For Aerial stem rot, begin applications prior to disease development and continue at 7-10 day intervals as needed. For White mold, begin foliar applications shortly after emergence or transplanting and continue on 7-10 day intervals if conditions are favorable for disease development. For Early blight and Late blight, Begin applications shortly after emergence or transplanting and continue on 5-7 day intervals as needed. Use Serifel in a rotational program or tank mix with other registered fungicides labeled for Early blight and Late blight control. For all other listed diseases, begin application prior to disease development and continue on 7-10 day intervals as needed. Mix and apply Serifel in sufficient water volume to ensure uniform dispersion in spray tank and thorough coverage of foliage and shoot tissue.
	Soil diseases Black rot of Sweet Potato* <i>(Ceratocystis fimbriata)</i> Black scurf <i>(Rhizoctonia solani)</i> Cavity spot <i>(Pythium spp.)</i> Fusarium wilt <i>(Fusarium spp.)</i> Phytophthora root rot <i>(Phytophthora spp.)</i> Pythium damping off <i>(Pythium spp.)</i> Rhizoctonia damping off <i>(Rhizoctonia solani)</i> Silver scurf* <i>(Helminthosporium spp.)</i> Verticillium wilt <i>(Verticillium spp.)</i>	4 – 16 oz/A (0.25 – 1.0 lb/A)	See Soil Application Instructions for In-furrow, Drench, Shank-In and Injected Applications. Apply a high enough water volume to thoroughly soak soil through the root zone.

*Not registered for use in California.

Crop-Specific Directions (continued)

Crop	Target Diseases	Product Use Rate per Application (oz or lb of Serifel®/Acre)	Application Directions
Tree nuts group Almond Beech nut Brazil nut Butternut Cashew Chestnut Chiquapin Coconut Filbert Hickory nut Pecan Pistachio Walnut (black and English)	Foliar diseases Alternaria blight (<i>Alternaria</i> spp.) Anthracnose (<i>Colletotrichum</i> spp.) Bacterial canker (<i>Pseudomonas syringae</i>) Botryosphaeria panicle and shoot blight* (<i>Botryosphaeria</i> spp.) Brown rot/Blossom blight (<i>Monilinia</i> spp.) Hull Rot (<i>Rhizopus</i> spp. and <i>Monilinia</i> spp.) Rust* (<i>Tranzschelia discolor</i>) Scab (<i>Cladosporium</i> spp.) Shot hole (<i>Wilsonomyces carpophilus</i>)	4 – 16 oz/A (0.25 – 1.0 lb/A)	For Anthracnose, Shot hole and Brown rot, begin applications prior to disease development when conditions are favorable for disease development and continue on 7-10 day intervals as needed. For Bacterial canker, begin applications prior to disease development and repeat at 7-10 day intervals as needed. Use the higher rate and shorter interval when disease pressure is high. For all other listed diseases, begin application prior to disease development and continue on 7 – 10 day intervals as needed. Mix and apply Serifel in sufficient water volume to ensure uniform dispersion in spray tank and thorough coverage of foliage and shoot tissue.

*Not registered for use in California.

Crop-Specific Directions (*continued*)

Crop	Target Disease	Product Use Rate per Application (oz or lb of Serifel®/Acre)	Application Directions
Wheat and Barley	Foliar diseases Head scab* (<i>Fusarium</i> spp.)	4 – 16 oz/A (0.25 – 1.0 lb/A)	For Head scab , apply Serifel at any time from one week prior to flowering to one week after flowering. Tank-mix Serifel with a fungicide labeled for head scab. Mix and apply Serifel in sufficient water volume to ensure uniform dispersion in spray tank and thorough coverage of foliage and shoot tissue.

*Not registered for use in California.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store in a cool, dry place until used. **DO NOT** store this product near food, feed, seed, fertilizers, or other pesticides.

Pesticide Disposal

To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

Container Handling

Nonrefillable Container. **DO NOT** reuse or refill this container. Clean container promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by state and local ordinances. If burned, stay out of smoke.

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing

Conditions of Sale and Warranty that may be varied only by agreement in writing signed by a duly authorized representative of BASF.

Serifel is a registered trademark of BASF.

© 2019 BASF Corporation
All rights reserved.

20190322.NVA 2017-04-492-0137
Based On: NVA 2017-04-492-0072

BASF Corporation 26 Davis Drive
Research Triangle Park, NC 27709



We create chemistry