

FIFRA Section 24(c) Special Local Need (SLN) Label



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Quintec®

**EPA 24(c) Special Local Need Registration SLN No. OR-180011
For Distribution and Use Only Within the State of Oregon**

**Expiration Date: This label is valid until December 31, 2023 or until otherwise amended,
withdrawn, canceled, or suspended.**

Spray Interval in Hops

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Quintec® fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Quintec according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Quintec.

Group	13	FUNGICIDE
--------------	-----------	------------------

Active Ingredient:

quinoxifen: 5,7-dichloro-4-(p-fluorophenoxy)
quinoline 22.58%
Other Ingredients 77.42%
Total 100.00%

Contains 2.08 lb of active ingredient per gallon.

Keep Out of Reach of Children

CAUTION

First Aid

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

Environmental Hazards

This product is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor runoff or drift from the target area. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

This product may contaminate water through drift of spray in wind. This product has a potential for runoff for several months or more after application. Poorly draining soils and soils with shallow watertables are more prone to produce runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

Directions for Use

Refer to product label for General Use Precautions, Mixing and Application instructions.

Maximum Application Rate: Do not apply more than 8.2 fl oz of Quintec® per acre per application or more than 33 fl oz of Quintec per acre per calendar year.

Minimum Spray Interval: 10 – 14 days. Use the shorter interval within the range for severe or threatening disease conditions.

Preharvest Interval: Do not apply within 21 days of harvest.

Quintec is in mode of action Group 13. Do not make more than four applications of Quintec per calendar year, nor more than two consecutive applications of Quintec before alternating to a fungicide that has a different mode of action and that is effective on local populations of the target powdery mildew pathogen. Alternate product should be applied within 14 days of applying Quintec.

Section 24c Registrant:
Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, IN 46268-1054 USA

®™Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

R136-075
Approved: 06/18/18
Initial printing.