RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

FIFRA 24(c) Special Local Need Label



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Lorsban® Advanced

EPA Reg. No. 62719-591

EPA 24(c) Special Local Need Registration SLN FL-090002 For Distribution and Use Only in the State of Florida

Control of Beet Armyworm, Fall Armyworm and Corn Earworm Larvae Infesting Sweet Corn

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Lorsban® Advanced insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Lorsban Advanced according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Lorsban Advanced.

Active Ingredient:

chlorpyrifos: O,O-diethyl-O-(3,5,6-trichloro-2-pyridinyl) phosphorothioate40.18%

Contains 3.755 lb of chlorpyrifos per gallon. Contains petroleum distillates.

Keep Out Of Reach Of Children WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Directions for Use

RESTRICTED USE PESTICIDE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Apply Lorsban Advanced at the rate of 1 pint per acre as a postemergence broadcast spray in sufficient water to ensure thorough coverage of plants using either aerial (fixed-wing or helicopter) or power-operated ground spray equipment. Use no less than 15 gallons of water per acre for ground spray equipment or at least 3 gallons of water per acre for aerial application. Control may be reduced at low spray volumes. Treat when field counts indicate that worms are or may become a problem.

Chemigation: Lorsban Advanced may be broadcast applied postemergence through sprinkler irrigation systems at specified application rates. See Chemigation Application section on product label for application instructions.

Specific Use Restrictions:

- Worker Restricted Entry Interval: Do not enter or allow worker entry into treated areas during the
 restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn. Refer to the
 Agricultural Use Requirements section on the product label for further information.
- Preharvest Interval: Do not apply within 7 days of harvest of ears.
- Do not make more than three foliar applications of Lorsban Advanced per season.
- Do not apply more than 3 lb ai chlorpyrifos per acre per season including all applications of chlorpyrifoscontaining products.
- Only one preplant application at the maximum of 1.3 lb ai chlorpyrifos per acre per season can be made along with the maximum of three foliar applications at 0.5 lb ai chlorpyrifos per acre.
- A minimum five-day reapplication interval is required for all repeat applications of Lorsban Advanced used as directed under this supplemental labeling.
- Maximum single application rate is 0.5 lb ai chlorpyrifos per acre.

Expiration date: This SLN labeling for Lorsban Advanced expires December 31, 2022 and is not valid for distribution or use after that date. Stocks of Lorsban Advanced purchased by growers on or prior to December 31, 2022 may continue to be used under this labeling until the supply of pesticide is exhausted.

Trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow

R368-130 Approved: 03/08/19 Replaces R368-115