

# CM

# CROP MIX ZIM w/B

## 5.0% S, 3% Zn, 2% Fe, 2% Mn, 0.2%B

### GENERAL INFORMATION

CROP MIX ZIM w/B is formulated for field, vegetable, row, fruit, tree, vine, ornamental and turf crops.

CROP MIX ZIM w/B may be applied through drip systems, aircraft, or via ground sprayers (high-volume or low-volume).

CROP MIX ZIM w/B may be applied alone or with fertilizers or pesticides. It can be applied either to the plant foliage or to the soil.

CROP MIX ZIM w/B is compatible with most pesticides and fertilizers. However, a compatibility test is recommended if the desired combination has not been used in the past.

### CONDITIONS OF SALE

Seller warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. Weather, crop conditions, timing and methods of applications, mixtures with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

SHAKE WELL BEFORE USE.

MANUFACTURED BY:

CUSTOM AGRICULTURAL FORMULATORS  
P.O. BOX 26104, FRESNO, CALIFORNIA 93729  
800-355-4457

### GUARANTEED ANALYSIS

Sulfur (S).....	5.0%
Boron(B).....	0.2%
Manganese (Mn).....	2.0%
Iron (Fe).....	2.0%
Zinc (Zn).....	3.0%

Derived from Zinc Sulfate, Ferrous Sulfate, Manganese Lignosulfonates and Boric Acid

Wt/Gal: 10.42 Lbs. @ 68°F 1249 g/ltr @ 20°C

### DIRECTIONS FOR USE

Always add this product to spray tank before adding a pesticide.

All Crops:

**Foliar Application:** Apply 2 to 4 quarts/acre as needed on a 10 - 14 day schedule. When rates of over 2 quarts are required, split applications of 2 quarts or less can be used.

Avoid foliar applications when plants are under stress due to lack of moisture.

### Soil Application:

For fertilizer maintenance apply 1-3 quarts per acre

For mild deficiency apply 2-6 quarts per acre

For high deficiency apply 1 - 3 gals per acre.

### Aerial Application:

Apply 1 quart per 5 to 10 gallons of water per acre.

### Drip Irrigation Application:

Apply the suggested rate of this product in multiple applications 10 - 14 days apart throughout the growing season of the crop.

### CAUTION:

This product contains Boron. Use of this product on any crops other than those recommended may result in serious injury to the crop(s).

**KEEP OUT OF  
REACH OF CHILDREN**



## WARNING

### PRECAUTIONARY STATEMENTS

**WARNING** Hazardous to humans and animals. Causes eye and skin irritations. When mixing or applying agricultural sprays, always wear spray goggles and impervious gloves plus protective clothing as recommended on pesticide labels.

### FIRST AID

If this product gets into the eyes, immediately wash the eyes with large amounts of water for at least 15 minutes, occasionally lifting the upper and lower lids. Obtain immediate medical treatment. If this product gets on the skin, immediately wash the contaminated area. Seek medical attention if skin irritation persists. If this product has been swallowed and the person is conscious, immediately give large volumes of water. Do not induce vomiting. Seek immediate medical attention.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Follow all recommended governmental procedures for disposal of pesticide spray solutions. Rinse container, empty rinsate and spray tank. Contact county agricultural commissioner for container disposal information.

Information regarding the contents and levels of metals in this product is available by calling (800) 355-4457.