

Classified for
"RESTRICTED USE"
in New York State
under 6NYCRR Part 326

FIFRA Section 24(c) Supplemental Label

RIMON® 0.83EC

Insecticide



**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF NEW YORK
FOR USE ONLY ON POTATOES IN NEW YORK STATE**

EPA REG. NO. 66222-35-400

EPA SLN NO. NY-130005

ACTIVE INGREDIENT:% BY WT.
Novaluron: 1-[3-chloro-4-(1,1,2-trifluoro-2-trifluoromethoxyethoxy)phenyl]-3-(2,6-difluorobenzoyl)urea*9.3%
OTHER INGREDIENTS:..... **90.7%**
Total..... 100.0%

*Contains 0.83 lbs. novaluron per gallon.

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

Pollinator Advisory: *In order to minimize the possibility of transient effects on honeybee brood development, do not use RIMON 0.83EC on blooming crops when bees are actively foraging. Do not allow product to drift into other flowering plants.*

FOR INSECT CONTROL IN POTATOES

Target Pests	Rates (Fl. Ozs. / A)	Application Instructions
Armyworms Colorado potato beetle European corn borer Foliage feeding caterpillars Loopers Potato tuberworm	6 to 12	Apply when the majority of the population is at egg hatch to the second instar. Use higher rates and higher spray volumes when larvae are large, or foliage canopy is tall or dense. Repeat applications as needed to protect new foliage growth, but not less than 7 days apart.
Do not apply to successive generations of Colorado potato beetle. Do not apply more than 24 oz. per acre per season. Do not apply within 14 days of harvest.		

continued on nex page

USE INFORMATION

- All applications in potatoes must include a 300-foot vegetative buffer strip between the edge of treated areas and bodies of water such as lakes, reservoirs, rivers, permanent streams, natural ponds, marshes or estuaries to decrease runoff.
- Apply required dosage by conventional ground sprayer equipment capable of delivering sufficient water to obtain thorough, uniform coverage of the target crop. Orient spray equipment boom and nozzles in a manner to minimize boom height, to optimize coverage uniformity, maximize deposition, and reduce spray drift. Drop nozzles may be required to obtain uniform coverage against certain pests that develop down in the canopy. Use a minimum of 10 gallons per acre in potatoes and vegetables. Higher gallonages will provide better coverage and performance. Use hollow cone, disc-core hollow cone or twin jet fan nozzles suitable for insecticide spraying.
- For ground boom applications, apply with nozzle height no more than 4 feet above the ground or crop canopy and when wind speed is 10 mph or less at the application site as measured by an anemometer. Use medium or coarser spray according to ASAE 572 definition for standard nozzles or VMD for spinning atomizer nozzles.

RESTRICTIONS

- Do not apply more than 3 applications per crop per season.
 - Do not apply within 14 days of harvest.
 - Apply by ground equipment only. Do not apply by air or through irrigation systems (chemigation).
 - Do not apply when wind speed favors drift beyond the area intended for treatment.
 - Do not allow RIMON to drift on grapes as leaf spotting may occur.
-

SLN Registrant:
Makhteshim-Agan of North America, Inc.
3120 Highwoods Blvd., Suite 100
Raleigh, NC 27604

Distributor:
Chemtura Corporation
199 Benson Road
Middlebury, CT 06749

Rimon is a registered trademark of Makhteshim Chemical Works Ltd.