



Bayer CropScience LP
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

Autumn™ Super 51 WDG Herbicide
EPA Reg. No. 264-1134

For Reduced Use Rates in Fall Applications

PRODUCT BULLETIN

FOR DISTRIBUTION AND USE ONLY IN THE STATES OF ALABAMA, ARIZONA, ARKANSAS, COLORADO, CONNECTICUT, DELAWARE, GEORGIA, IDAHO, ILLINOIS, INDIANA, IOWA, KANSAS, KENTUCKY, LOUISIANA, MAINE, MARYLAND, MASSACHUSETTS, MICHIGAN, MINNESOTA, MISSISSIPPI, MISSOURI, MONTANA, NEBRASKA, NEW JERSEY, NORTH CAROLINA, NORTH DAKOTA, OHIO, OKLAHOMA, OREGON, PENNSYLVANIA, RHODE ISLAND, SOUTH CAROLINA, SOUTH DAKOTA, TENNESSEE, UTAH, VERMONT, VIRGINIA, WASHINGTON, WEST VIRGINIA, WISCONSIN, WYOMING

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. This product bulletin should be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Autumn Super 51 WDG Herbicide may be applied at use rates ranging from 0.3 to 0.5 oz./A, depending upon geographic location, application timing and soil pH.

USE RATES

Time of Year	Geographical Use Location	Application Timing	Autumn Super 51 WDG Herbicide Use Rate (Oz/A)	
			Soil pH 6.8 or less	Soil pH 6.8 to 8.0 ¹
Fall	Areas North of Interstate 70	After Fall Harvest, Prior to Nov.1, Before ground freeze	0.5	0.3
		After Fall Harvest, After Nov. 1, Before ground freeze	0.3	0.3
	Areas South of Interstate 70	After Fall Harvest, Before ground freeze	0.5	0.3

¹DO NOT apply to soils with pH > 8.0

USE PRECAUTIONS

1. Autumn Super 51 WDG Herbicide will not provide season-long pre-emergence control of annual grass and broadleaf weeds. The length of residual weed control is generally rate dependent with greater residual control with higher use rates.
2. Application of an ALS-containing herbicide during the growing season following an Autumn Super 51 WDG herbicide application may increase the injury potential to soybeans when planted as the next rotational crop. In situations where extreme dry and cold conditions follow a Fall Autumn Super application, it is recommended to plant a STS soybean cultivar if one is using additional Spring-applied ALS-containing herbicides in soybeans.

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577
For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

*As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937)*