
SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300
SCP 953A-S10 0222

CYPRODINIL	GROUP	9	FUNGICIDE
FLUDIOXONIL	GROUP	12	FUNGICIDE

Switch® 62.5WG

Fungicide

This supplemental label expires on 02/04/2025 and must not be used or distributed after this date.

Active Ingredients:

Cyprodinil*	37.5%
Fludioxonil**	25.0%
Other Ingredients:	37.5%
Total:	100.0%

*CAS No. 121552-61-2

**CAS No. 131341-86-1

Switch 62.5WG is a water-dispersible granule containing 37.5% cyprodinil and 25% fludioxonil.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

EPA Reg. No. 100-953

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using Switch 62.5WG as permitted according to this Supplemental Labeling, read and follow all applicable directions, restrictions, and precautions on the EPA-registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and/or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of federal law to use this product in a manner inconsistent with its labeling.

syngenta

DIRECTIONS FOR USE

Crop	Disease	Product Rate oz/Acre	Use Directions
Brassica Head and Stem Vegetable Crop group 5-16** Broccoli Brussels sprouts Cabbage Cauliflower And cultivars and/or hybrids of these.	Powdery Mildew <i>(Erysiphe polygoni)</i>	10-12*	Begin applications prior to or at the onset of disease and repeat applications on a 7- to 10-day interval if conditions remain favorable for disease development.
	Alternaria leaf blight <i>(Alternaria spp.)</i> Suppression: Cercospora leaf spot <i>(Cercospora brassicicola)</i>	11-14*	Resistance Management: After 2 applications of Switch 62.5WG, alternate with another fungicide with a different mode of action for 2 applications.
*10 oz product/A contains 0.23 lb cyprodinil/A and 0.16 lb fludioxonil/A. *11 oz product/A contains 0.26 lb cyprodinil/A and 0.17 lb fludioxonil/A. *12 oz product/A contains 0.28 lb cyprodinil/A and 0.19 lb fludioxonil/A. *14 oz product/A contains 0.33 lb cyprodinil/A and 0.22 lb fludioxonil/A.			
**Complete List of Brassica Head and Stem Vegetables Crop group 5-16: Broccoli; Brussels sprouts; Cabbage; Cabbage, Chinese (napa); Cauliflower; cultivars, varieties, and hybrids of these commodities,			
Application Instructions: Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 10 gallons/A spray volume by air. Make no more than two applications by air. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			
Specific Use Restrictions <ol style="list-style-type: none"> Maximum Single Application Rate: Do not exceed the maximum rate listed in the table. Do not apply more than 4 applications per year at the highest rate. Minimum Application Interval: 7 days Make no more than two applications by air. Do not apply more than 56 oz/A of Switch 62.5WG per year (1.3 lb cyprodinil and 0.9 lb fludioxonil). Do not apply more than 1.3 lb ai/A of cyprodinil-containing products and 0.9 lb ai/A of fludioxonil-containing products per year. Do not apply within 7 days of harvest (7-day PHI). 			

Crop	Disease	Product Rate oz/Acre	Use Directions
Brassica Leafy Greens Subgroup 4-16B** Arugula Chinese cabbage Bok Choy Collards Kale Mustard greens Turnip greens And cultivars and/or hybrids of these. See separate instructions for watercress.	Powdery Mildew (<i>Erysiphe polygoni</i>) Alternaria leaf blight (<i>Alternaria</i> spp.) Suppression: Cercospora leaf spot (<i>Cercospora brassicicola</i>)	10-12* 11-14*	Begin applications prior to or at the onset of disease and repeat applications on a 7- to 10-day interval if conditions remain favorable for disease development. Resistance Management: After 2 applications of Switch 62.5WG, alternate with another fungicide with a different mode of action for 2 applications.
<div>*11 oz product/A contains 0.26 lb cyprodinil/A and 0.17 lb fludioxonil/A. *14 oz product/A contains 0.33 lb cyprodinil/A and 0.22 lb fludioxonil/A.</div>			
**Complete list of Brassica Leafy Greens Vegetable subgroup 4-16B: Arugula; broccoli, Chinese; broccoli raab; cabbage, abyssinian; cabbage, seakale; cabbage, Chinese, bok choy; collards; cress, garden; cress, upland; hanover salad; kale; maca, leaves; mizuna; mustard greens; radish, leaves; rape greens; rocket, wild; shepherd's purse; turnip greens; cultivars, varieties, and hybrids of these commodities.			
Application Instructions: Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 10 gallons/A spray volume by air. Make no more than two applications by air. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.			
Specific Use Restrictions <div>1) Maximum Single Application Rate: Do not exceed the maximum rate listed in the table. 2) Do not apply more than 4 applications per year at the highest rate. 3) Minimum Application Interval: 7 days 4) Make no more than two applications by air. 5) Do not use roots of treated turnips for food or feed. Only turnip varieties harvested for their leaves may be treated. 6) Do not apply more than 56 oz/A of Switch 62.5WG per year (1.3 lb cyprodinil and 0.9 lb fludioxonil). 7) Do not apply more than 1.3 lb ai/A of cyprodinil-containing products and 0.9 lb ai/A of fludioxonil-containing products per year. 8) Do not apply within 7 days of harvest (7-day PHI).</div>			

Crop	Disease	Product Rate oz/Acre	Use Directions
Citrus, Crop Subgroup 10-10B** Lemon Lime	Alternaria Stem End Rot (<i>A. citri</i>) Anthracnose (<i>Colletotrichum gloeosporioides</i>) Blue Mold (<i>Penicillium italicum</i>) Green Mold (<i>Penicillium digitatum</i>)	11-14*	Make one application near harvest to prevent post-harvest fruit rot. The application may be made up to and including the day of harvest.
*11 oz product/A contains 0.26 lb cyprodinil/A and 0.17 lb fludioxonil/A. *14 oz product/A contains 0.33 lb cyprodinil/A and 0.22 lb fludioxonil/A.			
**Complete List of Citrus Crop Group 10-10B: Australian desert lime; Australian finger-lime; Australian round lime; Brown River finger lime; kumquat; lemon; lime; mount white lime; New Guinea wild lime; Russell River lime; sweet lime; Tahiti lime; cultivars, varieties, and/or hybrids of these.			
Specific Use Restrictions 1) Maximum Single Application Rate: Do not exceed the maximum rate listed in the table. 2) Do not apply more than 1 application per year at the highest rate. 3) Application may be made by ground only. 4) Do not apply more than 14 oz/A of Switch 62.5WG per year (0.33 lb cyprodinil and 0.22 lb fludioxonil). 5) Do not apply more than 0.33 lb ai/A of cyprodinil-containing products and 0.22 lb ai/A of fludioxonil-containing products per year. 6) May be applied on the day of harvest (0-day PHI). 7) Do not exceed one application per year.			

Crop	Disease	Product Rate oz/Acre	Use Directions
Kohlrabi	Powdery Mildew (<i>Erysiphe polygoni</i>)	10-12*	Begin applications prior to or at the onset of disease and repeat applications on a 7- to 10-day interval if conditions remain favorable for disease development.
	Alternaria leaf blight (<i>Alternaria</i> spp.) Suppression: Cercospora leaf spot (<i>Cercospora brassicicola</i>)	11-14*	Resistance Management: After 2 applications of Switch 62.5WG, alternate with another fungicide with a different mode of action for 2 applications.
<p>*10 oz product/A contains 0.23 lb cyprodinil/A and 0.16 lb fludioxonil/A. *11 oz product/A contains 0.26 lb cyprodinil/A and 0.17 lb fludioxonil/A. *12 oz product/A contains 0.28 lb cyprodinil/A and 0.19 lb fludioxonil/A. *14 oz product/A contains 0.33 lb cyprodinil/A and 0.22 lb fludioxonil/A.</p>			
<p>Application Instructions: Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 10 gallons/A spray volume by air. Make no more than two applications by air. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.</p>			
<p>Specific Use Restrictions</p> <ol style="list-style-type: none"> 1) Maximum Single Application Rate: Do not exceed the maximum rate listed in the table. 2) Do not apply more than 4 applications per year at the highest rate. 3) Minimum Application Interval: 7 days 4) Make no more than two applications by air. 5) Do not apply more than 56 oz/A of Switch 62.5WG per year (1.3 lb cyprodinil and 0.9 lb fludioxonil). 6) Do not apply more than 1.3 lb ai/A of cyprodinil-containing products and 0.9 lb ai/A of fludioxonil-containing products per year. 7) Do not apply within 7 days of harvest (7-day PHI). 			

Crop	Disease	Product Rate oz/Acre	Use Directions
<p>Leafy Greens, Crop Subgroup 4-16A**, except Parsley</p> <p>Leaf petiole vegetables Crop Subgroup 22B***,</p> <p>Celtuce, Fennel, Florence, fresh leaves and stalk Celery Lettuce, head and leaf Spinach</p> <p>And cultivars and/or hybrids of these.</p>	<p>Alternaria leaf spot (<i>Alternaria</i> spp.)</p> <p>Septoria leaf spot (<i>Septoria lactucae</i>)</p> <p>Gray mold (<i>Botrytis cinerea</i>)</p> <p>Sclerotinia rot (<i>Sclerotinia</i> spp.)</p> <p>Basal rot (<i>Phoma exigua</i>)</p> <p>Suppression: Powdery mildew (<i>Erysiphe cichoracearum</i>)</p>	<p>11-14*</p>	<p>Begin applications prior to or at the onset of disease and repeat applications on a 7- to 10-day interval if conditions remain favorable for disease development.</p> <p>For control of Sclerotinia, make the first application at thinning and again two weeks later.</p> <p>Resistance Management: After 2 applications of Switch 62.5WG, alternate with another fungicide with a different mode of action for 2 applications.</p>
<p>*11 oz product/A contains 0.26 lb cyprodinil/A and 0.17 lb fludioxonil/A.</p>			
<p>*14 oz product/A contains 0.33 lb cyprodinil/A and 0.22 lb fludioxonil/A.</p>			
<p>**Complete List of Leafy Greens Crop subgroup 4-16A includes: Amaranth, Chinese; amaranth, leafy; aster, Indian; blackjack; cat's whiskers; cham-chwi; cham-na-mul; chervil, fresh leaves; chipilin; chrysanthemum, garland; cilantro, fresh leaves; corn salad; cosmos; dandelion, leaves; dang-gwi, leaves; dillweed; dock; dol-nam-mul; ebolo; endive; escarole; fameflower; feather cockscomb; good king henry; huauzontle; jute, leaves; lettuce, bitter; lettuce, head; lettuce, leaf; orach; plantain, buckhorn; primrose, English; purslane, garden; purslane, winter; radicchio; spinach; spinach, Malabar; spinach, New Zealand; spinach, tanier; Swiss chard; violet, Chinese, leaves; cultivars, varieties, and hybrids of these commodities</p> <p>***Leaf petiole vegetables Crop subgroup 22B includes: Cardoon; celery; celery, Chinese; fuki; rhubarb; udo; zuiki; cultivars, varieties, and hybrids of these commodities</p>			
<p>Application Instructions: Application may be made by ground, air, or chemigation. Good coverage is essential for good disease control. Use a minimum of 10 gallons/A spray volume by air. Make no more than two applications by air. For chemigation, apply in 0.1-0.25 inches/A of water. Chemigation with excessive water may lead to a decrease in efficacy.</p>			
<p>Specific Use Restrictions</p> <ol style="list-style-type: none"> 1) Maximum Single Application Rate: Do not exceed the maximum rate listed in the table. 2) Do not apply more than 4 applications per year at the highest rate. 3) Minimum Application Interval: 7 days 4) Make no more than two applications by air. 5) Do not apply more than 56 oz/A of Switch 62.5WG per year (1.3 lb cyprodinil and 0.9 lb fludioxonil). 6) Do not apply more than 1.3 lb ai/A of cyprodinil-containing products and 0.9 lb ai/A of fludioxonil-containing products per year. 7) May be applied on the day of harvest (0-day PHI). 			

Crop	Disease	Product Rate oz/Acre	Use Directions
Sugar Apple (Not for use in California)	Anthracnose (<i>Colletotrichum</i> spp.) Botrytis fruit rot (<i>Botrytis</i> spp.) Alternaria fruit rot (<i>Alternaria</i> spp.) Leaf and fruit spot (<i>Alternaria</i> spp.)	14*	Begin applications prior to disease development. Continue applications throughout the season on a 7-day interval, following resistance management guidelines. Resistance Management: After 2 applications of Switch 62.5WG, alternate with another fungicide with a different mode of action for 2 applications.
*14 oz product/A contains 0.33 lb cyprodinil/A and 0.22 lb fludioxonil/A.			
Application Instructions: Apply specified rate per acre as a directed foliar spray. Apply by ground using a minimum of 75 gallons/A spray volume. An adjuvant may be added at specified rates.			
Specific Use Restrictions 1) Do not apply via aerial application 2) Do not apply via irrigation, including chemigation. 3) Maximum Single Application Rate: Do not exceed the maximum rate listed in the table. 4) Do not apply more than 4 applications per year at the highest rate. 5) Minimum Application Interval: 7 days 6) Do not apply more than 56 oz/A of Switch 62.5WG per year (1.3 lb cyprodinil and 0.9 lb fludioxonil). 7) Do not apply more than 1.32 lb ai/A of cyprodinil-containing products and 0.88 lb ai/A of fludioxonil-containing products per year. 8) Do not use an organo-silicone adjuvant. 9) May be applied on the day of harvest (0-day PHI).			

Crop	Disease	Product Rate oz/Acre	Use Directions
Tropical and Subtropical, Small fruit, inedible peel subgroup 24A** Lychee Longan Spanish lime And cultivars and/or hybrids of these	Botrytis fruit rot (<i>Botrytis</i> spp.) Alternaria fruit rot (<i>Alternaria</i> spp.) Anthracnose (<i>Colletotrichum</i> spp.) Bipolaris bud and fruit rot (<i>Bipolaris</i> spp.)	11-14*	Make the first application during early bloom and repeat on 7- to 10-day intervals if conditions remain favorable for disease development. Resistance Management: After 2 applications of Switch 62.5WG, alternate with another fungicide with a different mode of action for 2 applications.
*11 oz product/A contains 0.26 lb cyprodinil/A and 0.17 lb fludioxonil/A. *14 oz product/A contains 0.33 lb cyprodinil/A and 0.22 lb fludioxonil/A.			
**Complete List of Subgroup 24A Crops: Aisen; bael fruit; Burmese grape; cat's-eyes; inga; longan; Lychee; madras-thorn; manduro; matisia; mesquite; mongongo, fruit; pawpaw, small-flower; satinleaf; Sierra Leone-tamarind; Spanish lime; velvet tamarind; wampi; white star apple; cultivars, varieties, and hybrids of these commodities.			
Application Instructions: Application may be made by ground or air. Good coverage is essential for good disease control. Use a minimum of 20 gallons/A spray volume by air. Make no more than two applications by air.			
Specific Use Restrictions 1) Maximum Single Application Rate: Do not exceed the maximum rate listed in the table. 2) Do not apply more than 4 applications per year at the highest rate. 3) Minimum Application Interval: 7 days 4) Make no more than two applications by air. 5) Do not apply more than 56 oz/A of Switch 62.5WG per year (1.3 lb cyprodinil and 0.9 lb fludioxonil). 6) Do not apply more than 1.3 lb ai/A of cyprodinil-containing products and 0.9 lb ai/A of fludioxonil-containing products per year. 7) Do not apply more than a maximum total of 4 applications (air plus ground) per year. 8) May be applied on the day of harvest (0-day PHI).			

Switch® 62.5WG and the Syngenta logo are trademarks of a Syngenta Group Company

©2022 Syngenta