SASOUTHERN DOAG CALCIUM NITRATE

DIRECTIONS FOR USE:

VEGETABLES and ANNUAL FLOWERS

As a supplemental (side dressing) fertilizer application. Apply 1 pound per 100 feet of row (1 cup per 50 feet of row or 1 level teaspoonful per plant). Apply to root growing area. Do not place within 2 inches of stalk or crown of plants. Do not allow contact with wet foliage. A second application two weeks after the first is suggested.

Calcium Nitrate may be used as a foliar spray for the addition of supplemental nitrogen and calcium. Apply as a full cover spray using 5 pounds per 100 gallons of spray. Applications within 1 week of harvest may adversely affect carrying quality of harvested produce.



ON APPLES AND OTHER FRUIT CROPS

Ground application broadcast under the tree. Use at rate of 1/2 pound for each year of tree growth to a maximum of 8 pounds per tree. Apply in February or apply in split applications in February and October.

Calcium Nitrate may be used as a foliar spray as an aid in the reduction of "bitter pit" and "cork spot" on apples. Apply with cover sprays every other week until 2 or 3 weeks before harvest using 3 pounds per hundred gallons of spray.

Calcium Nitrate is compatible with most insecticides and fungicides used on vegetables, floral crops and fruit crops. Do not use in combination with sulfates.

CAUTION: KEEP OUT OF REACH OF CHILDREN May be harmful if swallowed. Avoid contact with eyes.

Do not store with combustible materials or nitrogenous tankage. In case of fire, flood with water. Sweep up spilled material. Close partially used bags. Store in a safe, dry place.

GUARANTEED ANALYSIS:



Southern Agricultural Insecticides, Inc.

F672

Palmetto, FL 34220

.12

P.O. Box 218, Palmetto, FL 34220

Hendersonville, NC 28793

Boone, NC 28607

Net Weight: 5 lbs. (2.27 kg.)

S A SOUTHERN 5 D AG_®

CALCIUM NITRATE