



## 2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA

# DUPONT™ EXIREL® INSECT CONTROL

EPA REG. NO. 352-859

## FOR THE CONTROL OF DIAPREPES ROOT WEEVIL ADULTS IN CITRUS IN THE STATE OF FLORIDA

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Do not apply this product in any way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

EXIREL® must be used only in accordance with directions on this bulletin or in separately published DuPont labeling.

EXIREL® is recommended for control of Diaprepes root weevil adults in citrus.

Apply EXIREL® at 13.5 to 20.5 fluid ounces per acre as a foliar spray. For best performance, use of an adjuvant such as horticultural spray oil at 1 gallon per 100 gallons of spray volume (1% v/v) is recommended (read adjuvant, compatibility and tank mix and crop safety sections of main label). Target applications according to local thresholds. When targeting Diaprepes root weevil and other labeled pests such as Asian citrus psyllid, citrus leafminer and cotton aphid time applications to early flush and use a minimum EXIREL® rate of 16 fluid ounces per acre plus an adjuvant as above.

Spray Volume: Thorough coverage is essential to achieve best results. Select a spray volume appropriate for the size of trees or plants and density of foliage.

Where higher spray volumes are used, apply a higher EXIREL® rate in the specified rate range. For best results, apply 100-150 gallons of water per acre when using commercial airblast equipment. Do not apply less than 30 gallons of water per acre when using commercial airblast equipment.

Read and follow the protection of pollinators section of the main label when using during bloom.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with such non-labeled use.

### RESTRICTIONS

Do not apply a total of more than 0.4 lb ai/A of CYAZYPYR® or cyantraniliprole containing products per year.

Re-entry interval is 12 hours.

Last application (days to harvest) = 1 day.

(Replaces R-1910 092414 10-01-14)

#### IMPORTANT NOTICE

**BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.**

It is a violation of federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This bulletin should be in the possession of the user at the time of application.

**Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.**

Copies of DuPont product labels are available by calling 1-888-6-DUPONT  
or at <http://cropprotection.dupont.com/>

©2017 E. I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Road, Wilmington, DE 19805 U.S.A.