



Bayer Environmental Science

Bayer Environmental Science
P.O. Box 12014
2 T.W. Alexander Drive
Research Triangle Park, North Carolina 27709
1-866-99BAYER (1-866-992-2937)

Premise® 2 Insecticide

EPA Reg. No. 432-1331

For control of Outdoor/perimeter pests

PRODUCT BULLETIN

FOR DISTRIBUTION AND USE ONLY IN THE STATES OF ALABAMA, ARIZONA, ARKANSAS, CALIFORNIA, COLORADO, CONNECTICUT, DELAWARE, DISTRICT OF COLUMBIA, FLORIDA, GEORGIA, ILLINOIS, INDIANA, IOWA, KANSAS, KENTUCKY, MAINE, MARYLAND, MASSACHUSETTS, MICHIGAN, MINNESOTA, MISSISSIPPI, MISSOURI, MONTANA, NEBRASKA, NEVADA, NEW JERSEY, NORTH CAROLINA, NORTH DAKOTA, OHIO, OKLAHOMA, OREGON, PENNSYLVANIA, PUERTO RICO, RHODE ISLAND, SOUTH CAROLINA, SOUTH DAKOTA, TENNESSEE, UTAH, VERMONT, VIRGINIA, WASHINGTON, WEST VIRGINIA, WISCONSIN AND WYOMING

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. It is a violation of federal law to use this product in a manner inconsistent with its label. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its label.

APPLICATION FOR OUTSIDE/PERIMETER PESTS

Outdoor perimeter pests including ants, boxelder bugs, crickets, earwigs, elm leaf beetle, firebrats, ground beetles, millipedes, moths, pillbugs, silverfish, sowbugs, springtails, stink bugs (including brown marmorated stink bug and globular stink bug [kudzu bug]).

Apply to surfaces on buildings, porches, patios and other structures, around doors and windows, eaves and attic vents, utility entry points, soffit areas and other exterior openings (including foundation cracks or drilled holes) where these pests enter the structure or where they have been seen or found, or can find shelter. Apply low pressure (do not exceed 50 psi at the nozzle tip) mist or spray.

To prevent infestation, treat the outside of the structure, to include walls, around doors, windows, and soffits where pests are active or may enter or hide. Treat the soil, turf or other ground covering adjacent to the structure (in a band up to 10 feet wide) where pests are seen, have been found, or can find shelter.

Apply directly to insects/pests that have congregated around doors, windows, and similar areas where they may enter the structure. To help reduce infestations, treat cracks and crevices, attics, and other winter hiding places located inside buildings.

High volume applications using power sprayers: You may adjust the rate of product to ensure that 0.27 fl. ounces (8ml.) of this product is applied per 1000 square feet of treated surface. For example; mixing 4.5 fluid ounces of this product in a 50 gallon tank, while using an application volume of 3.0 gallons finished spray solution per 1000 square feet, will apply 0.27 fluid ounces of this product per 1000 square feet. The table below gives examples of end-use dilutions for typical power spraying application volumes.

DILUTION RATE CHART		
Amount of Product fluid ounces (milliliter)	Application Volume gallons water	Concentration (% AI) in spray solution
0.27 (8)	1	0.05%
	2	0.025%
	3	0.017%
	4	0.013%
	5	0.01%

Treatments may be applied monthly to maintain control.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information, Call 1-866-99BAYER (1-866-992-2937)

*As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*