



**DuPont Crop  
Protection**

**FIFRA Section 24(c)  
SPECIAL LOCAL NEED**

**DUPONT™ VERIMARK®  
INSECT CONTROL**

For Distribution and Use Only Within the State of Oregon

**DUPONT™ VERIMARK® INSECT CONTROL**

EPA Reg. No. 352-860

EPA SLN No. OR-170004

**FOR SUPPRESSION OF CABBAGE MAGGOT IN TURNIP-ROOTED  
PARSLEY, RADISH, ORIENTAL RADISH (DAIKON), RUTABAGA AND  
TURNIP GROWN FOR SEED PRODUCTION**

This label is valid until December 31, 2022 or until otherwise amended, withdrawn, canceled, or suspended.

**Active Ingredient**

Cyantraniliprole  
3-bromo-1-(3-chloro-2-pyridinyl)-N-[4-cyano-2-methyl-6-[(methylamino)  
carbonyl]phenyl]-1H-pyrazole-5-carboxamide

Other Ingredients

**TOTAL**

**By Weight**

18.66%

81.34%

**100.00%**

**KEEP OUT OF REACH OF CHILDREN**

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to aquatic invertebrates and oysters. Do not apply directly to water. Drift and runoff may be hazardous to aquatic organisms in water adjacent to use sites. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are foraging the treatment area.

**Surface Water Advisory-**

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several weeks after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of cyantraniliprole from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

**Ground Water Advisory-**

This chemical has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling. This Special Local Need (SLN) label and the federally registered main product label for Verimark® must be in possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this SLN label and the main EPA-registered product label. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

© 2017 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Rd., Wilmington, DE 19805 U.S.A

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks with uses not specified by the label or DuPont 24(c) instructions.

DuPont™ VERIMARK® Insect Control is recommended for the suppression of Cabbage Maggot in turnip-rooted parsley, radish, oriental radish (daikon), rutabaga and turnip grown for seed production only.

Apply VERIMARK® at 10.0 to 13.5 fluid ounces per acre as a soil at planting application (in-furrow spray, surface band, soil shank injection, transplant tray drench by growers or commercial transplant producers no earlier than 72 hours prior to planting in the field, transplant water treatment or hill drench). VERIMARK® must be applied uniformly in the root zone to ensure effective control. Surface band application requires a sufficient amount of water post-application to ensure the treatment is moved into the root zone. See the SOIL APPLICATION section of the federal label for additional guidance, also see the rate conversion chart for application rate per 1000 linear feet.

Adjusting the pH of the application solution: The pH of the application solution should be between pH 4 and 6. (See "Acidification of Application Solution" section of the federal label.)

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with uses not specified by the label or DuPont 24(c) instructions.

## **RESTRICTIONS**

Do not make ground applications within 25' of lakes, rivers, reservoirs, permanent streams, marshes, natural ponds, estuaries or coastal areas. Do not cultivate within 25' of these aquatic areas to allow growth of a vegetative filter strip.

Do not apply more than 0.4 lbs ai/A of CYAZYPYR® or cyantraniliprole containing products per year whether applications are made to the soil or foliarly.

The field re-entry interval is 4 hours.

For use only on specified crops grown for seed.

## **IMPORTANT NOTICE:**

**BEFORE BUYING OR USING DUPONT™ VERIMARK®, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS IN THIS SECTION 24(c) LABEL AND ON THE EPA-REGISTERED LABEL.**

This label and the federal label for this product must be in the possession of the user at the time of pesticide application.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

Read the Limitation of Warranty and Liability on the product label before buying or using this product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the product label.