
SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 1202A-S8 1120

MEFENOXAM	GROUP	4	FUNGICIDE
-----------	-------	---	-----------

Ridomil Gold® SL

Fungicide

This supplemental label expires on 11/09/2023 and must not be used or distributed after this date.

Active Ingredient:

Mefenoxam*	45.3%
------------------	-------

<i>Other Ingredients:</i>	54.7%
---------------------------	-------

Total:	100.0%
--------	--------

*CAS Nos. 70630-17-0 and 69516-34-3

Ridomil Gold® SL is formulated as a soluble concentrate and contains 4 lb active ingredient per gallon.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

EPA Reg. No. 100-1202

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using Ridomil Gold SL as permitted according to this Supplemental Labeling, read and follow all applicable directions, restrictions, and precautions on the EPA-registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and/or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

syngenta

Tree Nuts, Crop Group 14-12

Crops (including all cultivars, varieties, and/or hybrids)			
African nut-tree Almond Beechnut Brazil nut Brazilian Pine Bunya Bur Oak Butternut Cajou nut Candlenut	Cashew Chestnut Chinquapin Coconut Coquito nut Dika nut Ginkgo Guiana Chestnut Hazelnut (Filbert) Heartnut	Hickory nut Japanese horse-chestnut Macadamia Nut Mongongo nut Monkey pot Monkey puzzle nut Okari nut Pachira nut Peach palm nut	Pecan Pequi Nut Pili Nut Pine nut Pistachio Sapucaia nut Tropical almond Walnut, black Walnut, English Yellowhorn
Target Disease	Rate pt/A (lb ai)	Application Timing	Use Directions
Collar rot Crown rot Root rot (<i>Phytophthora</i> spp.)	4.0 (2.0) or 1.5 fl oz/1,000 sq ft	2 weeks after planting (new plantings) or in the spring before growth begins (established plantings). Additional applications may be made at 2- to 3-month intervals, depending on disease pressure. Apply before symptoms appear.	Soil spray (broadcast, band or irrigation): Apply to soil beneath the tree or apply through irrigation water (micro-sprinkler or drip) to cover the root zone. For intensive plantings (2-3 times the normal planting rate), apply on a per area basis (1,000 sq ft).
Resistance Management: <ul style="list-style-type: none"> Refer to Section 3.2 			
Precaution: <ul style="list-style-type: none"> Ridomil Gold SL will not revitalize trees showing moderate to severe disease symptoms. 			
USE RESTRICTIONS			
1) Refer to Section 6.1 for additional product use restrictions. 2) Maximum Single Application Rate: 4.0 pt/A (equivalent to 2.0 lb ai/A mefenoxam) 3) Minimum Application Interval: 60 days 4) Maximum Annual Application Rate: 12.0 pt/A/year (equivalent to 6.0 lb ai/A/year mefenoxam) a) DO NOT exceed 6.0 lb ai/A/year of soil-applied mefenoxam- and metalaxyl-containing products. 5) DO NOT apply more than 3 applications per year. 6) DO NOT concentrate spray around tree trunks. 7) DO NOT apply to trees under stress. 8) In CA, DO NOT apply to newly planted trees within 45 days of planting. On some varieties, chlorosis may occur on leaf margins. 9) DO NOT graze livestock in treated areas. 10) DO NOT graze or feed cover crops grown in treated orchards. 11) Pre-harvest Interval (PHI): 30 days			

Ridomil Gold® SL and the Syngenta logo are trademarks of a Syngenta Group Company.