

PEEL HERE TO OPEN ►



INDOXACARB GROUP 22 INSECTICIDE

Advion[®] Insect

Granular bait

syngenta[®]

Insecticide

A granular bait for indoor and outdoor control of cockroaches, crickets (including mole crickets), nuisance ants, and listed crawling insect pests in and around residential, commercial, public, and institutional buildings.

Active Ingredient:

Indoxacarb

(S)-methyl 7-chloro-2,5-dihydro-2-[[[(methoxycarbonyl)
[4(trifluoromethoxy) phenyl]amino]carbonyl]indeno

[1,2-e][1,3,4]oxadiazine-4a-(3H)carboxylate 0.22%

Other Ingredients 99.78%

Total: 100.00%

**KEEP OUT OF REACH OF CHILDREN
CAUTION/PRECAUCIÓN**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

EPA Reg. No. 100-1483 EPA Est. 39578-TX-1

**SCP 1483B-L1 1219
4116307**

PRODUCT ID
66718

1 pound
Net Contents



FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment

HOTLINE NUMBER

For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-888-8372

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Avoid contact with skin, eyes or clothing.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Shoes plus socks.
- Long-sleeved shirt and long pants are required when using hand-held application equipment.

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment rinsewater or rinsate. Do not apply where/when conditions could favor runoff. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas.

Do not apply to any impervious surfaces which may contact or lead directly to surface water, storm drains, or urban runoff conveyance systems (gutters). Sweeping any product that lands on an impervious surface back onto pervious surfaces will prevent runoff to water bodies or drainage systems. Cover, incorporate, or clean up granules that are spilled.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product. Syngenta will not be responsible for losses or damages resulting from use of this product in any manner not specifically stated on this label or other labels or bulletins published by Syngenta. User assumes all risks associated with such non-specified use.

PRODUCT INFORMATION

Advion Insect Granular Bait is a ready-to-use granular bait that contains the active ingredient indoxacarb. When applied according to label use directions, Advion Insect Granular Bait controls the wide range of insect pests listed in Table 1.

RESISTANCE MANAGEMENT

INDOXACARB	GROUP	22	INSECTICIDE
------------	-------	----	-------------

For resistance management, Advion Insect Granular Bait contains a Group 22A insecticide. Any insect population may contain individuals naturally resistant to Advion Insect Granular Bait and other Group 22A insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same treated areas. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance, take the following steps:

- Rotate the use of Advion Insect Granular Bait or other Group 22A insecticides with different groups that control the same pests.
- Adopt an integrated pest management program for insecticide use that includes scouting, uses historical information related to pesticide use, record keeping, and which considers cultural, biological, and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.

Contact your local Syngenta representative, retailer, or extension specialist for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area. For further information or to report suspected resistance contact Syngenta at 1-866-SYNGENT(A) (866-796-4368). Syngenta encourages responsible product stewardship to ensure effective long-term control of the insects on this label.

USE RESTRICTIONS

- Not for use on turfgrass that is being grown for seed or on commercial sod farms.
- Not for use in nurseries and greenhouses.
- DO NOT allow livestock or domestic animals to consume the bait or graze in treated areas.
- DO NOT apply indoors to areas accessible to children or pets.
- DO NOT allow Advion Insect Granular Bait to contact any food or feed.
- Do not apply this product through any type of irrigation system.
- DO NOT exceed 4.6 pounds of Advion Insect Granular Bait per 1000 sq ft per year or 3.2 pounds of Advion Insect Granular Bait per 100 linear ft applied in a 7 ft band (equal to 0.44 lb active ingredient per acre per year).
- Do not formulate this product into other end-use products.

OUTDOOR APPLICATION

Advion Insect Granular Bait contains the active ingredient indoxacarb in a formulation that is highly attractive to the pests listed in the following table.

Table 1. Treatment Rates and Pests Controlled

Inside and around residential dwellings, business and office complexes, shopping complexes, multi-family residential complexes, institutional buildings, airports, cemeteries, parks, playgrounds, schools, including lawns and landscapes, sports and athletic fields, and other similar landscape turf areas			
Pest Controlled	lb product/ 1,000 sq ft	lb product/ 100 linear ft	lb AI/ acre
ants, including tawny crazy ants (except red imported fire ants, harvester ants, carpenter ants and pharaoh ants) earwigs field crickets house crickets millipedes ³ mole cricket adults and nymphs (<i>Scapteriscus</i> spp.) ⁴ silverfish	0.38 to 4.6 ¹	0.27 to 3.2 ¹	0.037 to 0.44
cockroaches	1.15 to 4.6 ²	0.81 to 3.2 ²	0.11 to 0.44

¹ equivalent to 0.61 to 7.36 oz per 100 sq ft

² equivalent to 1.84 to 7.36 oz per 100 sq ft

³ not approved for use in California

⁴ minimum rate is 1.15 lb per 1000 sq ft in California

Structural Perimeter-Band Application

To control pests listed in the table above except mole crickets, apply to ground area in a band up to 7 feet wide around and adjacent to the structure. For mole cricket control, see section on Limited Area Applications.

Landscape Boundary Application

To control pests listed in the table above, apply to fence lines, ornamental gardens, pavement crevices, planter areas, and soil, vegetation, and mulched areas.

Limited Area Applications (Spot Treatment) for Listed Pests in Lawns and Landscapes

Apply Advion Insect Granular Bait at a rate of 0.38 to 4.6 lb product per 1000 sq ft as a limited area treatment. Repeat application after 7 days, if necessary. To control adult mole crickets before they lay eggs, Advion Insect Granular Bait should be applied in the spring. To control mole crickets nymphs and newly emerged adults, apply Advion Insect Granular Bait in the late summer and fall.

For best results, apply in late afternoon to areas of turfgrass that are infested with mole crickets, show evidence of mole cricket damage or have a history of mole cricket infestation. If the soil is not moist, then it is important to irrigate before application to bring the mole crickets to the soil surface where exposure to Advion Insect Granular Bait will be maximized.

Bait Station - Apply bait using commercial bait stations suitable for granular insect bait. Place 2 to 4 oz of bait in each bait station depending on the bait station design and size. The number of bait stations placed in the treated area will depend on the size of the area and the level of the infestation. Monitor the bait stations, and replenish bait granules as needed.

Table 2. Outdoor Perimeter-Band Application Rates

Application Rate lb/1,000 sq ft	Band Width (feet)	lb of Advion® Insect Granular Bait for Perimeter Applications Linear Distance (feet) Around Structure			
		100	150	200	250
0.38	5	0.19	0.29	0.38	0.48
0.38	7	0.27	0.40	0.53	0.67
0.5	5	0.25	0.38	0.50	0.63
0.5	7	0.35	0.53	0.70	0.88
1.15	5	0.58	0.86	1.15	1.44
1.15	7	0.81	1.21	1.61	2.01
2.3	5	1.15	1.73	2.30	2.88
2.3	7	1.61	2.42	3.22	4.03
3.0	5	1.50	2.25	3.00	3.75
3.0	7	2.10	3.15	4.20	5.25
3.5	5	1.75	2.63	3.50	4.38
3.5	7	2.45	3.68	4.90	6.13
4.0	5	2.00	3.00	4.00	5.00
4.0	7	2.80	4.20	5.60	7.00
4.6	5	2.30	3.45	4.60	5.75
4.6	7	3.22	4.83	6.44	8.05

Application Rate lb/1,000 sq ft	Band Width (feet)	lb of Advion® Insect Granular Bait for Perimeter Applications Linear Distance (feet) Around Structure			
		300	400	500	1000
0.38	5	0.57	0.76	0.95	1.90
0.38	7	0.80	1.06	1.33	2.66
0.5	5	0.75	1.00	1.25	2.50
0.5	7	1.05	1.40	1.75	3.50
1.15	5	1.73	2.30	2.88	5.75
1.15	7	2.42	3.22	4.03	8.05
2.3	5	3.45	4.60	5.75	11.50
2.3	7	4.83	6.44	8.05	16.10
3.0	5	4.50	6.00	7.50	15.00
3.0	7	6.30	8.40	10.50	21.00
3.5	5	5.25	7.00	8.75	17.50
3.5	7	7.35	9.80	12.25	24.50
4.0	5	6.00	8.00	10.00	20.00
4.0	7	8.40	11.20	14.00	28.00
4.6	5	6.90	9.20	11.50	23.00
4.6	7	9.66	12.88	16.10	32.20

INDOOR APPLICATION

Advion Insect Granular Bait may be used to control the pests listed in Table 3 in crawl spaces, finished and unfinished basements and attics of residential, industrial and commercial buildings or structures. Advion Insect Granular Bait will kill cockroaches that consume the bait granules when placed in attics, crawl spaces and basements of residential, industrial and commercial buildings or structures, but this type of application alone is not intended to provide complete control of an indoor cockroach infestation. Monitor the treated area, and reapply bait granules only as needed.

For use in food or feed handling establishments, see section below.

Crack & Crevice and Void Application – Apply bait into cracks and crevices or voids at a rate up to 2 oz per 10 linear feet using a bulb applicator or other equipment for applying granular bait into cracks & crevices or voids.

Scatter Application - Apply bait as scatter treatment at rate of 0.38 lb to 4.6 lb of product per 1,000 sq ft. The minimum application rate for cockroaches is 1.15 lb product per 1,000 sq ft (equivalent to 1.84 oz per 100 sq ft).

Perimeter Band Treatment - Apply bait along walls in a band approximately 6 to 12 inches wide at rate of 2 oz bait granules per 10 linear feet.

Bait Station - Apply bait using commercial bait stations suitable for granular insect bait. Place 2 to 4 oz of bait in each bait station depending on the bait station design and size. The number of bait stations placed in the treated area will depend on the size of the area and the level of the infestation.

Table 3. Indoor Treatment Methods and Pests Controlled

Pests Controlled	Treatment Method	Treatment Location	Treatment Rate
ants, including tawny crazy ants (except fire ants, harvester ants, carpenter ants and pharaoh ants) cockroaches ¹ earwigs field crickets house crickets millipedes ² silverfish	Crack & Crevice, Void treatment	• Indoors	up to 2 oz per 10 linear feet
	Scatter baiting	• Crawl spaces • Attics (uninhabited)	up to 7.36 oz per 100 square feet
	Perimeter band	• Crawl spaces • Unfinished basements • Attics (uninhabited)	up to 2 oz per 10 linear feet in a band 6 to 12 inches wide
	Bait station	• Crawl spaces • Unfinished basements • Finished basements • Attics	2 to 4 oz per bait station (number of bait stations will depend on the level of infestation)

¹ minimum application rate for cockroaches is 1.84 oz per 100 sq ft

² not approved for use in California

DO NOT apply to areas accessible to children and pets.

FOOD- AND FEED-HANDLING ESTABLISHMENTS

(Food-handling establishments are places other than private residences in which food is held, processed, prepared or served.)

Applications in Non-food/Non-feed Areas

Advion Insect Granular Bait may be used to control listed pests in residential structures and the non-food areas of commercial, industrial, public and institutional buildings/structures, including restaurants, warehouses, food processing plants, supermarkets, hospitals, nursing homes, motels, hotels, schools, daycares, laboratories, computer facilities, aircraft, buses, boats/ships, trains, and pet shops.

Non-food and non-feed areas include areas such as garbage rooms, lavatories, floor drains (to sewers), entries, and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after bottling or canning). Refer to section below for use directions and restrictions when making applications in food- and feed-handling areas of Food and Feed Handling Establishments.

Applications in Food/Feed Areas

Food- and feed-handling areas include areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing edible waste storage, and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas are also considered a food/feed area when food is exposed and facility is in operation.

Apply between different elements of construction, between floor and equipment, openings to voids and hollow spaces in walls, bases and equipment or using bait stations.

Specific Restrictions for Food- and Feed-Handling Establishments

- **DO NOT** apply to areas or surfaces where food/feed, food utensils or food-processing surfaces may come into contact and become contaminated.
- **DO NOT** apply to areas that are routinely washed, such tops of tables, food/feed preparation and prepared food-holding surfaces.

STORAGE AND DISPOSAL

Prohibitions

Do not contaminate water, food or feed by storage and disposal.

Pesticide Storage

Store product in original container only in a location inaccessible to children and domestic animals/pets. Store in a cool, dry place and avoid excessive heat.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal

Non-refillable container. Do not reuse or refill this container. Completely empty plastic container, plastic bag or fiber sack into application equipment by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then offer for recycling, if available or dispose of plastic container, plastic bag or fiber sack in a sanitary landfill, or by other procedures approved by state and local authorities.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, LLC or Seller. To the extent permitted by applicable law, Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and, (2) Buyer and User assume the risk of any such use. **TO THE EXTENT PERMITTED BY**

APPLICABLE LAW, SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.

To the extent permitted by applicable law, in no event shall SYNGENTA be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

Advion®, the ALLIANCE FRAME
the SYNGENTA Logo
and the PURPOSE ICON
are Trademarks of a
Syngenta Group Company
©2019 Syngenta



For non-emergency
(e.g., current product information), call
Syngenta Crop Protection at 1-800-334-9481

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300