

LIQUID NICKEL 5%

A Formulation for the Correction of Nickel Deficiencies in Crops

GENERAL INFORMATION

CM LIQUID NICKEL 5% may be applied through drip systems, aircraft, or via ground sprayers (high-volume or low-volume).

CM LIQUID NICKEL 5% may be applied alone or with fertilizers or pesticides. It can be applied either to the plant foliage or to the soil.

CM LIQUID NICKEL 5% is compatible with most pesticides and fertilizers. However, a compatibility test is recommended if the desired combination has not been used in the past.

SHAKE WELL BEFORE USE.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Follow proper procedures for the disposal of spray solutions. Rinse and dispose of empty container according to approved local practices.

Information regarding the contents and levels of metals in this product is also available by calling 800-355-4457.

MANUFACTURED BY: CUSTOM AG FORMULATORS, INC PO BOX 26104 FRESNO, CA 93729 800-355-4457

DANGER

REACH OF CHILDREN

KEEP OUT OF

PRECAUTIONARY STATEMENTS

WARNING Hazardous to humans and animals. Causes eye and skin irritations. Ingestion may cause nausea. When mixing or applying agricultural sprays, always wear spray goggles and impervious gloves plus protective clothing as recommended on pesticide labels.

FIRST AID

If this product gets into the eyes, immediately wash the eyes with large amounts of water for at least 15 minutes, occasionally lifting the upper and lower lids. Obtain immediate medical treatment. If this product gets on the skin, immediately wash the contaminated area. Seek medical attention if skin irritation persists. If this product has been swallowed and the person is conscious, immediately give large volumes of water. Do not induce vomiting. Seek immediate medical attention.

DIRECTIONS FOR USE

Always add this product to spray tank after filling with water but before adding a pesticide or fertilizer. The addition of a non-ionic surfactant can be beneficial.

FOLIAR APPLICATIONS:

Pecan and Nut Crop:

First Application: Parachute stage.

Apply 1 to 1.5 pints/acre. 2 applications per season are recommended, With first application at parachute stage.

Second Applications: Apply Mid-July

Apply 1.5 to 2 pints/acre.

Avoid foliar applications of **CM LIQUID NICKEL 5%** when plants are under stress due to lack of moisture.

Peanuts: Apply 8 oz/acre in adequate water beginning 50 days after planting and repeat every 14-21 days apart throughout the growing season of the crop.

Soybeans:

Apply 10-16 oz/acre in adequate water beginning at R-1 and repeat in 2 -3 weeks.

Cotton:

Apply 8 oz./acre with growth regulator application to control vegetative growth.